



**Cover:** The hearing capabilities of diving birds are not well understood. Using positive reinforcement training, a great cormorant (*Phalacrocorax carbo*), was trained to voluntarily participate in psychophysical hearing research to help us better understand the in-air hearing capabilities of this species. The great cormorant is more sensitive to in-air sounds than previously believed and its hearing abilities are comparable to several other species of birds of similar size. This knowledge is important for our understanding of the hearing abilities of other species of sea birds. See the article by Maxwell et al. on page 496. Image provided by Alyssa Maxwell and licensed under a Creative Commons Attribution 4.0 International licence.

## RESEARCH ARTICLES

- 407** Investigation of feet functions of large ruminants with a decoupled model of equivalent mechanism  
**Zhang, Q., Xu, K. and Ding, X.**
- 415** Reprogramming towards totipotency is greatly facilitated by synergistic effects of small molecules  
**Miyamoto, K., Tajima, Y., Yoshida, K., Oikawa, M., Azuma, R., Allen, G. E., Tsujikawa, T., Tsukaguchi, T., Bradshaw, C. R., Jullien, J., Yamagata, K., Matsumoto, K., Anzai, M., Imai, H., Gurdon, J. B. and Yamada, M.**
- 425** Non-centralized and functionally localized nervous system of ophiuroids: evidence from topical anesthetic experiments  
**Matsuzaka, Y., Sato, E., Kano, T., Aonuma, H. and Ishiguro, A.**
- 439** tBRD-1 and tBRD-2 regulate expression of genes necessary for spermatid differentiation  
**Theofel, I., Bartkuhn, M., Boettger, T., Gärtner, S. M. K., Kreher, J., Brehm, A. and Rathke, C.**
- 449** A novel immunodeficient NOD.SCID-*rd1* mouse model of retinitis pigmentosa to investigate potential therapeutics and pathogenesis of retinal degeneration  
**Mishra, A., Das, B., Nath, M., Iyer, S., Kesarwani, A., Bhattacharjee, J., Arindkar, S., Sahay, P., Jain, K., Sahu, P., Sinha, P., Velpandian, T., Nagarajan, P. and Upadhyay, P.**
- 463** *Xenopus laevis* Kif18A is a highly processive kinesin required for meiotic spindle integrity  
**Möckel, M. M., Heim, A., Tischer, T. and Mayer, T. U.**
- 471** Metabolic rate and hypoxia tolerance are affected by group interactions and sex in the fruit fly (*Drosophila melanogaster*): new data and a literature survey  
**Burggren, W., Souder, B. M. and Ho, D. H.**
- 481** *Mycobacterium tuberculosis* Cpn60.2 (GroEL2) blocks macrophage apoptosis via interaction with mitochondrial mortalin  
**Joseph, S., Yuen, A., Singh, V. and Hmama, Z.**
- 489** Tanshinone IIA suppresses the progression of atherosclerosis by inhibiting the apoptosis of vascular smooth muscle cells and the proliferation and migration of macrophages induced by ox-LDL  
**Wang, B., Ge, Z., Cheng, Z. and Zhao, Z.**
- 496** In-air hearing of the great cormorant (*Phalacrocorax carbo*)  
**Maxwell, A., Hansen, K. A., Ortiz, S. T., Larsen, O. N., Siebert, U. and Wahlberg, M.**
- 503** Propulsive efficiency of frog swimming with different feet and swimming patterns  
**Jizhuang, F., Wei, Z., Bowen, Y. and Gangfeng, L.**
- 511** Depletion of *Tcf3* and *Lef1* maintains mouse embryonic stem cell self-renewal  
**Ye, S., Zhang, T., Tong, C., Zhou, X., He, K., Ban, Q., Liu, D. and Ying, Q.-L.**
- 518** Novel device for application of continuous mechanical tensile strain to mammalian cells  
**Wada, S., Kanzaki, H., Narimiya, T. and Nakamura, Y.**
- 525** Time-varying auditory gain control in response to double-pulse stimuli in harbour porpoises is not mediated by a stapedial reflex  
**Schröder, A. E. M., Beedholm, K. and Madsen, P. T.**