

RESEARCH ARTICLE

How does the snakehead *Channa argus* survive in air? The combined roles of the suprabranchial chamber and physiological regulations during aerial respiration

Ting Duan^{1,*}, Chenchen Shi^{1,*}, Jing Zhou², Xiao Lv¹, Yongli Li¹ and Yiping Luo^{1,‡}

ABSTRACT

This study aimed to test the hypothesis that the aerial survival of the northern snakehead is involved not only with suprabranchial chamber respiration but also with physiological regulations. The aerial survival time and oxygen consumption rate (VO_2) were determined in snakeheads with either normal or injured suprabranchial organs. Some hematological and biochemical parameters were assessed during aerial exposure. The results showed that resting VO_2 decreased when switching from water to air in both the control and the suprabranchial organ-injured fish, with decreases of 22.4% and 23.5%, respectively. Resting VO_2 in air was not different between the control and the suprabranchial organ-injured fish. The red blood cell (RBC) count and hemoglobin concentration showed no marked changes, while RBC size increased when exposed to air. The liver lactate concentration remained unchanged, and the white muscle lactate concentration decreased when switching from water to air. The blood ammonia concentration tended to increase during aerial respiration. These results suggest that the aerial survival of the snakehead is positively associated with a combination of factors, including respiration of suprabranchial organs and other accessory organs, depressed metabolic demands and increased oxygen transport, and negatively associated with the accumulation of blood ammonia but not anaerobic metabolism.

KEY WORDS: Respiration, *Channa argus*, Metabolic demand, Oxygen consumption, Anaerobic metabolism

INTRODUCTION

The northern snakehead (*Channa argus*) is a species of air-breathing fish that is widely distributed in East Asia. This fish has a peculiar accessory breathing organ, the suprabranchial chamber, by which the fish breathes air at the water's surface using a cough-like mechanism (Ishimatsu and Itazawa, 1981; Lefevre et al., 2014), making possible short-term survival out of water (Nagata and Nakata, 1988). The suprabranchial chamber of *Channa* possesses a richly vascularized wall for gas exchange. It communicates with the

buccopharyngeal cavity through a ventral opening guarded by a 'shutter' plate bone outgrowth of the first branchial arch, which can close the inhalant aperture of the suprabranchial chamber during expiration (Munshi, 1962). Even though many air-breathing fish maintain equivalent oxygen consumption in air and in water (Yoshiyama and Cech, 1994; Sayer, 2005), some air-breathing fish reduce their oxygen consumption rate (VO_2) in air (Garey, 1962; Martin et al., 2004; Lefevre et al., 2014). Our recent study has found that the northern snakehead can depress metabolic level in air (Li et al., 2017), implying that the snakehead may have mechanisms other than its suprabranchial chamber to meet metabolic demands and survive out of water.

In addition to ventilation, some other physiological regulations may be necessary for fish to breathe air (Sayer, 2005). Fish may improve their blood oxygen-carrying capacity to compensate for their impaired oxygen uptake and/or enhance anaerobic metabolism (which is indicated by lactic acid concentration) during aerial exposure (Farmer, 1979; Bennett, 1978; Graham, 1997). Whether these regulations occur in the northern snakehead during aerial exposure to compensate for air respiration of the suprabranchial organ remains unclear.

Short-term aerial survival has been reported in several *Channa* species, e.g., 8 h for *C. striata*, >27 h for *C. batrachus* (S. Chandra, Histopathology of respiratory organs of fishes, PhD thesis, Banaras Hindu University, 2001; Chandra and Banerjee, 2004), and 14–24 h for the northern snakehead, depending on ambient temperature and metabolic demand (Li et al., 2017). It has been suggested that air respiration of the snakehead via the cough-like ventilation mechanism is water dependent (Ishimatsu and Itazawa, 1981), and therefore that the snakehead would be unable to survive in terrestrial habitats for very long (Liem, 1984). An alternative mechanism for the aerial survival of fish could be ammonia poisoning. In water, the ammonia products of fish are excreted easily and diffuse rapidly (Mommensen and Walsh, 1992; Wood, 1993). Out of water, however, the ammonia excretion of the fish may be blocked, which could lead to an accumulation of ammonia in the fish body (Chew et al., 2003; Sayer, 2005). Therefore, it can be hypothesized that the limited aerial survival of the northern snakehead could be attributed to excess ammonia accumulation.

We assumed that the aerial survival of the snakehead is involved not only with suprabranchial chamber respiration but also with physiological regulations, such as depressed metabolic demands, enhanced blood oxygen-carrying capacity and enhanced anaerobic metabolism. The oxygen consumption of fish with normal or injured suprabranchial chambers was compared to test the contribution of the suprabranchial chamber to oxygen uptake. VO_2 , blood parameters, lactate content of liver and muscle, and plasma ammonia were also assessed to test the changes in oxygen-carrying capacity, anaerobic metabolism and ammonia

¹Key Laboratory of Freshwater Fish Reproduction and Development, Ministry of Education, School of Life Sciences, Southwest University, Chongqing 400715, China. ²Department of Clinical Medicine, Chongqing Medical and Pharmaceutical College, Chongqing 401331, China.

*These authors contributed equally to this work

‡Author for correspondence (luoguo@swu.edu.cn).

 Y.Luo, 0000-0003-3312-6403

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Received 21 August 2017; Accepted 15 January 2018

accumulation of the fish when switched from water-breathing to air-breathing states.

RESULTS

The VO_2 of both the control and suprabranchial organ-injured fish decreased when switching from water respiration to aerial respiration (Fig. 1). Resting VO_2 in air (VO_{2Air}) was significantly lower than resting VO_2 in water (VO_{2Water}), with decreases of 22.4% in the control and 23.5% in the injured fish (Table 1). Resting VO_{2Air} did not significantly differ between the control and injured fish. The aerial survival time of the injured fish was 19.0 h, nearly 4 h less than that of the control (22.9 h).

The oxygen-carrying capacity related parameters, including red blood cell (RBC) count (RBCC), hemoglobin concentration (Hb) and mean cellular hemoglobin content (MCH), of the fish did not change markedly when switching from water respiration to aerial respiration (Table 2). The RBC length (RBCL), width (RBCW) and area (RBCA) of the fish after several hours of aerial respiration were significantly higher than those of the fish after water respiration. The lactate content ranged from 2.6 to 3.1 $mg\ g^{-1}$ in the liver and from 1.4 to 1.9 $mg\ g^{-1}$ in the white muscle of the fish (Fig. 2). The liver lactate content did not significantly change, while the white muscle lactate content remained unchanged (at 3 or 6 h) or significantly decreased (at 1.5 and 12 h) when switching from water respiration to aerial respiration.

The blood ammonia concentration of the fish in water was 0.29 $mmol\ l^{-1}$ and tended to increase during aerial respiration (Fig. 3). After 12 h of aerial respiration, the blood ammonia concentration reached 0.50 $mmol\ l^{-1}$.

DISCUSSION

The suprabranchial chamber of the snakehead is an important accessory breathing organ that allows gas exchange in air (Ishimatsu and Itazawa, 1981; Glass et al., 1986). However, in our study, injury to the suprabranchial chamber did not change the VO_{2Air} of the northern snakehead, which suggests that organs other than the

suprabranchial chamber also contribute to aerial respiration. Consistently, previous histological studies have shown that the roof of the buccopharynx and the surface of the tongue of the snakehead are also vascularized structures (Ishimatsu et al., 1979; Ishimatsu and Itazawa, 1983a,b).

Consistent with our previous study (Li et al., 2017), the snakeheads showed lower VO_{2Air} than VO_{2Water} (Fig. 1). A similarly reduced VO_2 in air has also been observed in the Atlantic silverside, *Menidia menidia* (Garey, 1962; Martin et al., 2004). A reduced VO_2 in fish can be attributed to a lower capacity for gas exchange or a downregulated metabolic demand. It has been proposed that the ventilation of the snakehead is inefficient during air exposure (Ishimatsu and Itazawa, 1981; Li et al., 2017), which limits its terrestrial survival capacity (Liem, 1984). However, the reduced VO_2 of the snakehead in air might be due to different mechanisms, as the group with suprabranchial organ injuries maintained a VO_2 equal to that of the fish with intact organs, suggesting that this species has sufficient oxygen exchange capacity. Therefore, the reduced VO_2 of the snakehead in air can be ascribed to a downregulation of metabolic demand (Li et al., 2017). A lower metabolic rate is meaningful for fish in extreme environments, allowing them to save energy, to reduce end-product accumulation, and to survive for longer periods (Ginneken and Thillart, 2009).

RBCs and Hb are useful indicators of the blood oxygen-carrying capacity. Fish with a larger RBCC have greater blood oxygen-carrying capacity. For fish in water, the blood oxygen-carrying capacity may increase when faced with an oxygen shortage (Weber and Fago, 2004), and splenic erythrocyte release can increase RBCC and Hb (Randall, 1982; Murad et al., 1990; Pearson and Stevens, 1991). However, these hematological parameters of some air-breathing fish may not change during air exposure, as sufficient oxygen supply can be maintained by the air-breathing organ (Cruz et al., 2013). Similarly, in our study, the RBCC, Hb and MCH of the snakehead showed no marked changes in air compared to those in water, suggesting a sufficient oxygen supply. Interestingly, the RBC




Fig. 1. Oxygen consumption rate (VO_2) of northern snakeheads switched from water respiration to aerial respiration. $n=14$ for each group. Data are presented as the mean \pm s.e.m. The values with symbols (*, control group; †, suprabranchial organ group) were different from the resting VO_2 in water by a paired samples t -test ($P<0.05$). Filled circles with solid line, control group; open circles with dashed line, suprabranchial organ group.

Table 1. Water and aerial respiration parameters of northern snakeheads

Treatment	Control	Suprabranchial organ injured	<i>t</i>	<i>P</i>
Number	14	14		
Body mass (g)	3.20±0.04	3.19±0.04	0.290	0.774
Resting VO _{2Water} (mg O ₂ kg ⁻¹ h ⁻¹)	266.4±10.3	263.1±9.0	0.240	0.812
Resting VO _{2Air} (mg O ₂ kg ⁻¹ h ⁻¹)	202.2±4.9	198.6±4.4	0.551	0.587
Change in resting VO ₂ from water to air (%)	22.4±4.1	23.5±2.9	-0.232	0.819
Survival time (h)	22.9±0.5 ^a	19.0±0.8 ^b	4.281	0.000224

Data are presented as the mean±s.e.m. ^{a,b}Values with a superscript of a different letter are significantly different (*P*<0.05).

Table 2. Hematological parameters of northern snakeheads switched from water respiration to aerial respiration

Time (h)	0	1.5	3	6	12	<i>F</i>	<i>P</i>
Number	10	9	10	10	9		
Body mass (g)	8.39±0.1	8.02±0.1	8.31±0.1	8.45±0.1	8.37±0.1	1.761	0.154
RBCL (μm)	8.84±0.08 ^a	8.67±0.12 ^a	9.33±0.18 ^b	9.22±0.15 ^b	9.36±0.12 ^b	5.326	0.001
RBCW (μm)	6.01±0.10 ^a	6.52±0.16 ^{bc}	6.80±0.17 ^b	6.36±0.14 ^{a,c}	6.37±0.10 ^{ac}	4.534	0.004
RBCA (μm ²)	41.76±0.68 ^a	44.45±1.46 ^{a,d}	49.93±1.86 ^b	46.09±1.28 ^{c,d}	46.87±1.16 ^{b,c,d}	5.146	0.002
RBCC (10 ⁹ ml ⁻¹)	2.31±0.12	2.41±0.31	2.45±0.31	2.56±0.17	2.74±0.17	0.486	0.746
Hb (mg ml ⁻¹)	60.44±2.00	56.56±3.66	57.28±4.25	53.65±3.37	59.58±2.63	0.678	0.611
Mean cellular Hb (pg)	26.73±1.51	24.89±1.98	24.82±1.93	21.98±2.24	22.33±1.42	1.151	0.346

Data are presented as the mean±s.e.m. ^{a,b,c,d}Values with a superscript of a different letter are significantly different (*P*<0.05).

size increased when the fish were exposed to air, suggesting some extent of erythrocyte swelling, which can improve oxygen transport in fish (Wendelaar Bonga, 1997). Our results suggest that the aerial metabolic demand of snakeheads depends more on the regulation of RBC surface area than on the numbers of RBCs.

Another mechanism that allows fish to withstand environments with low oxygen availability is anaerobic metabolism (Hochachka et al., 1996). In our study, the anaerobic metabolism of the snakehead was not increased in air, as reflected by the unchanged liver lactate concentration and even decreased white muscle lactate concentration during aerial respiration (Fig. 2). This suggests that anaerobic metabolism has a minor contribution to the aerial survival of the northern snakehead. Unaffected liver and white muscle lactate has also been observed in another air-breathing fish, *Pterygoplichthys anisitsi* (Cruz et al., 2013).

Ammonia is the most toxic of the respiratory gases of fishes (Ip et al., 2001). When exposed to air, some fishes can maintain unchanged blood ammonia concentrations by reducing the rates of proteolysis and amino acid catabolism, by continuing to excrete

ammonia, or by converting ammonia to less toxic forms, allowing them to survive in air for a long time; these species include the giant mudskipper (*Periophthalmodon schlosseri*), the swamp eel (*Monopterusuchia*) and the African lungfish (*Protopterus annectens*) (Anderson, 2001; Ip et al., 2004; Chew et al., 2005). In contrast, accumulation of blood ammonia generally appears in fishes with reduced ammonia excretion (Chew et al., 2003), which may limit their ability to survive in air. Similarly, in our study, the blood ammonia of the northern snakehead accumulated during aerial respiration, which may partly explain its limited aerial survival time (23 h).

In conclusion, when northern snakeheads are exposed to air, their survival is positively associated with a combination of factors, including the respiration of suprabranchial chambers and other accessory respiratory organs, depressed metabolic demands and increased oxygen transport, but is negatively associated with the accumulation of blood ammonia. The regulation of anaerobic metabolism has a minor contribution to aerial survival of the northern snakehead.




Fig. 2. Lactate concentrations in white muscle and liver of northern snakeheads switched from water respiration to aerial respiration. *n*=10, 9, 10, 10 and 9 for the fish post-aerial respiration for 0 (in water), 1.5, 3, 6 and 12 h, respectively. Data are presented as the mean±s.e.m. The values of each organ with asterisks were different from values of the fish in water (0 h) by an independent samples *t*-test (*P*<0.05). Filled column, white muscle; open column, liver.




Fig. 3. Blood ammonia concentrations of northern snakeheads switched from water respiration to aerial respiration. $n=10, 9, 10, 10,$ and 9 for the fish post-aerial respiration for 0 (in water), 1.5, 3, 6 and 12 h, respectively. Data are presented as the mean \pm s.e.m. The values with asterisks were different from values of the fish in water (0 h) by an independent samples *t*-test ($P<0.05$).

MATERIALS AND METHODS

The snakeheads were obtained from Huashan Hatchery, Guangdong Province, China. The fish were acclimated in a rearing system at the Fisheries Science Institute of Southwest University. The water temperature was $25\pm 1^\circ\text{C}$, and the photoperiod was 12 h light: 12 h dark. The oxygen concentration was $>6\text{ mg l}^{-1}$, and the ammonia concentration was $<0.015\text{ mg l}^{-1}$. Animal handling and experiments were approved by School of Life Sciences, Southwest University (LS-SWU-1612) and were conducted according to the requirements of Environment and Housing Facilities for Laboratory Animals of China (GB/T14925-2001).

To test the change in VO_2 due to suprabranchial organ injury, the snakeheads were divided into two groups: a control group ($n=14$) and an injured group ($n=14$). The fish were fasted for 24 h, and the body mass was weighed ($\sim 3\text{ g}$ each). Then, the fish were individually transferred into respiratory chambers and adapted overnight before determination of $\text{VO}_{2\text{Water}}$ and $\text{VO}_{2\text{Air}}$. The structure of the respiratory chamber and the methods for VO_2 determination were described by Li et al. (2017). Briefly, the chamber (30 ml) can be switched from a flow-through water phase to a closed air phase by regulating the inlet and outlet valves. For water respiration, the dissolved oxygen concentration ($\text{mg O}_2\text{ L}^{-1}$) was measured at the outlet by a fiber optic sensor system (MircroX TX3, PresSens Precision Sensing GmbH, Regensburg, Germany), and the water flow rate (L h^{-1}) was measured by collecting water over a given time. The $\text{VO}_{2\text{Water}}$ ($\text{mg O}_2\text{ h}^{-1}$) was calculated by multiplying the water flow rate by the difference in dissolved oxygen concentration between the fish chamber and the control chamber. The dissolved oxygen in the outlet water was maintained at $>70\%$ saturation concentration to avoid causing hypoxic stress. For $\text{VO}_{2\text{Air}}$ determination, the water in the chamber was discharged, and a near-saturated humidity condition was maintained in the chamber. $\text{VO}_{2\text{Air}}$ was determined in an intermittent flow pattern. The initial oxygen partial pressure of the air was measured, and the chamber was sealed for 90 min; then, the final oxygen level was measured. The change in air volume due to air breathing was determined from the reading on a 1-ml syringe connected to the inlet. Atmospheric pressure was recorded during each measurement. Then, the air in the chamber was refreshed for 30 min, and the next determination loop was started. The oxygen concentration in the air ($\text{mg O}_2\text{ l}^{-1}$) was obtained from the values of oxygen partial pressure, temperature and atmospheric pressure. $\text{VO}_{2\text{Air}}$ ($\text{mg O}_2\text{ h}^{-1}$) was calculated as $\text{VO}_{2\text{Air}}=(\text{O}_{2i}\times V_i-\text{O}_{2f}\times V_f)/t$, where O_{2i} and O_{2f} are the initial and final oxygen concentrations of the air ($\text{mg O}_2\text{ l}^{-1}$), respectively; V_i and V_f are the initial and final air volume, respectively; and t is the breathing time (h). The volume of the fish body was estimated by assuming a body density of 1 kg l^{-1} . One chamber without fish was set as a control for either water or air respiration. A mass-specific value of VO_2 was obtained by dividing by body mass (kg). $\text{VO}_{2\text{Water}}$ was measured for both groups at 1-h intervals for 8 h, and the mean value was taken as the resting $\text{VO}_{2\text{Water}}$. For the injured group, we sheared the ventral and

posterior walls of the suprabranchial chamber using surgical scissors. After 30 min of adaptation, the $\text{VO}_{2\text{Air}}$ of the two groups was determined at 2-h intervals for 24 h. The mean values were taken as the resting $\text{VO}_{2\text{Air}}$.

Another series of experiments was conducted to test the changes in hematological and biochemical parameters during aerial exposure. Snakeheads ($\sim 8\text{ g}$) were individually placed in moist beakers (1 l) in the air at $25\pm 1^\circ\text{C}$. After 0 (control), 1.5, 3, 6 or 12 h of aerial exposure, fish were anaesthetized by immersion in MS222 solution (tricaine methanesulfonate) for 2–3 min (Luo et al., 2013). A blood sample was taken by syringe from the caudal vein and divided into four parts to determine the RBCC, RBC size, Hb and blood ammonia concentration. The liver and a piece of white muscle below the dorsal fin were sampled on ice using surgical scissors for the determination of lactate concentration. The RBCC was determined with a Neubauer counter, and the RBCL and RBCW of 50 randomly selected RBCs were determined under a digital light microscope (EV5680, Aigo Company, Beijing, China) linked to a computer. The RBCA was calculated using the formula: $\text{RBCA}=\text{RBCL}\times\text{RBCW}\times\pi/4$ (Gao et al., 2007). Hb was determined by the alkalized hemoglobin method (Dacie and Lewis, 1984). The MCH was calculated by the ratio of Hb to RBCC. The blood ammonia concentration was determined with a blood ammonia determination kit (Jiancheng Biotech Co., Ltd, Nanjing, China). White muscle and liver were each mechanically homogenized and diluted 10-fold with ice-cold saline solution (0.65%), and the contents of lactate were measured using a lactic acid assay kit (Jiancheng Biotech Co., Ltd, Nanjing, China).

Data analysis

Data were calculated using Microsoft Excel 2007 and statistical analyses were performed using IBM SPSS 11.5. The independent samples *t*-test was used to compare resting VO_2 , change in VO_2 when switching from water to air, and survival time between the control and injured groups. The paired samples *t*-test was used to compare the resting $\text{VO}_{2\text{Water}}$ with the $\text{VO}_{2\text{Air}}$ at each time point. One-way ANOVA followed by a multiple comparisons test (LSD) was used to test the changes in the blood parameters along time points during aerial exposure. The independent samples *t*-test was used to compare lactate content and ammonia concentration at each time point during aerial respiration with those during water respiration. Data are presented as the mean \pm s.e.m. Differences were considered statistically significant when $P<0.05$.

Acknowledgements

We thank Bo Zhang for assisting with fish collection and Ilouise Bradford for help with English writing.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: Y. Luo; Methodology: T.D., C.S., J.Z., X.L., Y. Li, Y. Luo; Software: T.D., C.S., J.Z., Y. Li, Y. Luo; Validation: J.Z., X.L., Y. Luo; Formal analysis: T.D.; Investigation: T.D., C.S., J.Z., X.L., Y. Li, Y. Luo; Resources: C.S.; Data curation: T.D., C.S., Y. Luo; Writing - original draft: T.D., C.S., Y. Luo; Writing - review & editing: T.D., C.S., J.Z., X.L., Y. Li, Y. Luo; Supervision: Y. Luo; Project administration: Y. Luo; Funding acquisition: Y. Luo.

Funding

This work was supported by the National Natural Science Foundation of China [31672287].

Data availability

Data for this study have been deposited in FigShare (Duan, 2018), doi:10.6084/m9.figshare.5844477.

References

- Anderson, P. M.** (2001). Urea and glutamine synthesis: environmental influences on nitrogen excretion. In *Fish Physiology, Nitrogen Excretion* (ed. P. A. Wright and P. M. Anderson), pp. 239-277. New York: Academic Press.
- Bennett, A. F.** (1978). Activity metabolism of the lower vertebrates. *Ann. Rev. Physiol.* **40**, 447-469.
- Chandra, S. and Banerjee, T. K.** (2004). Histopathological analysis of the respiratory organs of *Channa striata* subjected to air exposure. *Vet. Arhiv.* **74**, 37-52.
- Chew, S. F., Wong, M. Y., Tam, W. L. and Ip, Y. K.** (2003). The snakehead *Channa asiatica* accumulates alanine during aerial exposure, but is in capable of sustaining locomotory activities on land through partial amino acid catabolism. *J. Exp. Biol.* **206**, 639-704.
- Chew, S. F., Wilson, J. M., Ip, Y. K. and Randall, D. J.** (2005). Nitrogenous excretion and defense against ammonia toxicity. In *Fish Physiology, the Physiology of Tropical Fishes* (ed. A. Val, V. Almeda-Val and D. J. Randall), pp. 307-395. New York: Academic Press.
- Cruz, A. L., Silva, H. R., Lundstedt, L. M., Schwantes, A. R., Moraes, G., Klein, W. and Fernandes, M. N.** (2013). Air-breathing behavior and physiological responses to hypoxia and air exposure in the air-breathing loricariid fish, *Pterygoplichthys anisitsi*. *Fish Physiol. Biochem.* **39**, 243-256.
- Dacie, J. V. and Lewis, S. N.** (1984). *Practical Hematology*, 5th edn. Edinburgh: Churchill Livingstone.
- Farmer, M.** (1979). The transition from water to air breathing: effects of CO₂ on hemoglobin function. *Comp. Biochem. Physiol. A.* **62**, 109-114.
- Gao, Z. X., Wang, W. M., Abbas, K., Zhou, X. Y., Yang, Y., Diana, J. S., Wang, H. P., Wang, H. L., Li, Y. and Sun, Y. H.** (2007). Haematological characterization of loach *Misgurnus anguillicaudatus*: comparison among diploid, triploid and tetraploid specimens. *Comp. Biochem. Physiol. A.* **147**, 1001-1008.
- Garey, W. F.** (1962). Cardiac responses of fishes in asphyxic environments. *Biol. Bull. -US.* **122**, 362-368.
- Ginneken, V. and van den Thillart, G.** (2009). Metabolic depression in fish measured by direct calorimetry: a review. *Thermochim. Acta.* **483**, 1-7.
- Glass, M. L., Ishimatsu, A. and Johansen, K.** (1986). Responses of aerial ventilation to hypoxia and hypercapnia in *Channa argus*, an air-breathing fish. *J. Comp. Physiol. B.* **156**, 425-430.
- Graham, J. B.** (1997). The families of air-breathing fishes. In *Air-Breathing Fishes: Evolution, Diversity and Adaptation* (ed. J. B. Graham), pp. 57-59. New York: Academic Press.
- Hochachka, P. W., Buck, L. T., Doll, C. J. and Land, S. C.** (1996). Unifying theory of hypoxia tolerance: molecular metabolic defense and rescue mechanisms for surviving oxygen lack. *PNAS* **93**, 9493-9498.
- Ip, Y. K., Chew, S. F. and Randall, D. J.** (2001). Ammonia toxicity, tolerance and excretion. In *Fish Physiology, Nitrogen Excretion* (ed. P. A. Wright and P. M. Anderson), pp. 109-148. New York: New Academic Press.
- Ip, Y. K., Chew, S. F. and Randall, D. J.** (2004). Five tropical air-breathing fishes, six different strategies to defend against ammonia toxicity on land. *Physiol. Biochem. Zool.* **77**, 768-782.
- Ishimatsu, A. and Itazawa, Y.** (1981). Ventilation of the air-breathing organ in the snakehead *Channa argus*. *Jap. J. Ichthyol.* **28**, 276-282.
- Ishimatsu, A. and Itazawa, Y.** (1983a). Blood oxygen levels and acid-base status following air exposure in an air-breathing fish, *Channa argus*: The role of air ventilation. *Comp. Biochem. Physiol.* **74**, 787-793.
- Ishimatsu, A. and Itazawa, Y.** (1983b). Difference in blood oxygen levels in the outflow vessels of the heart of an air-breathing fish, *Channa argus*: do separate blood streams exist in a teleostean heart? *J. Comp. Physiol.* **149**, 435-440.
- Ishimatsu, A., Itazawa, Y. and Takeda, T.** (1979). On the circulatory systems of the snakeheads *Channa maculata* and *C. argus* with reference to bimodal breathing. *Jap. J. Ichthyol.* **26**, 167-180.
- Lefevre, S., Wang, T. and Jensen, A., Cong, N. V., Huong, D. T. T., Phuong, N. T. and Bayley, M.** (2014). Air-breathing fishes in aquaculture. What can we learn from physiology? *J. Fish Biol.* **84**, 705-731.
- Li, Y. L., Lv, X., Zhou, J., Shi, C. C., Duan, T. and Luo, Y. P.** (2017). Does air-breathing meet metabolic demands of the juvenile snakehead, *Channa argus*, in multiple conditions. *Biol. Open.* **6**, 642-664.
- Liem, K. F.** (1984). The muscular basis of aquatic and aerial ventilation in the air-breathing teleost fish *Channa*. *J. Exp. Biol.* **113**, 1-18.
- Luo, Y. P., Wang, W., Zhang, Y. R., Huang, Q. D. and Lim, D.** (2013). Effects of starvation on the excess post-exercise oxygen consumption of juvenile Nile tilapia (*Oreochromis niloticus*). *Mar. Freshw. Behav. Phys.* **45**, 333-342.
- Martin, K. L. M., Van Winkle, R. C., Draais, J. E. and Lakisic, H.** (2004). Beach spawning fishes, terrestrial eggs, and air breathing. *Physiol. Biochem. Zool.* **77**, 750-759.
- Mommsen, T. P. and Walsh, P. J.** (1992). Biochemical and environmental perspectives on nitrogen metabolism in fishes. *Experientia.* **48**, 583-593.
- Munshi, J. S. D.** (1962). On the accessory respiratory organs of *Ophicephalus punctatus* (Bloch.) and *Ophicephalus striatus* (Bloch.). *Zool. J. Linn. Soc.* **44**, 616-626.
- Murad, A., Houston, A. H. and Samson, L.** (1990). Haematological response to reduced oxygen-carrying capacity, increased temperature and hypoxia in goldfish, *Carassius auratus* L. *J. Fish. Biol.* **36**, 289-305.
- Nagata, Y. and Nakata, Y.** (1988). Distribution of six species of bitterlings in a creek in Fukuoka Prefecture, Japan. *Jap. J. Ichthyol.* **35**, 320-331.
- Pearson, M. P. and Stevens, E. D.** (1991). Size and hematological impact of the splenic erythrocyte reservoir in rainbow trout, *Oncorhynchus mykiss*. *Fish Physiol. Biochem.* **9**, 39-50.
- Randall, D. J.** (1982). The control of respiration and circulation in fish during exercise and hypoxia. *J. Exp. Biol.* **100**, 275-288.
- Sayer, M. D. J.** (2005). Adaptations of amphibious fish for surviving life out of water. *Fish. Fish.* **6**, 186-211.
- Weber, R. E. and Fago, A.** (2004). Functional adaptation and its molecular basis in vertebrate hemoglobins, neuroglobins and cytoglobins. *Respir. Physiol. Neurobiol.* **144**, 141-159.
- Wendelaar Bonga, S. E.** (1997). The stress response in fish. *Physiol. Rev.* **77**, 591-625.
- Wood, C. M.** (1993). Ammonia and urea metabolism and excretion. In *The Physiology of Fishes* (ed. D. H. Evans), pp. 379-425. Boca Raton: CRC Press.
- Yoshiyama, R. M. and Cech, J. J.** (1994). Aerial respiration by rocky intertidal fishes of California and Oregon. *Copeia.* **1994**, 153-158.