

RESEARCH ARTICLE

Sustained exercise-trained juvenile black carp (*Mylopharyngodon piceus*) at a moderate water velocity exhibit improved aerobic swimming performance and increased postprandial metabolic responses

Xiuming Li^{1,2}, Yaoguang Zhang², Xiaojin Li¹, Hua Zheng¹, Jianglan Peng¹ and Shijian Fu^{1,*}

ABSTRACT

The objectives of this study were to examine whether sustained exercise training at four water velocities, i.e. nearly still water (control), 1 body length (BL) s⁻¹, 2 BL s⁻¹ and 4 BL s⁻¹, has effects on swimming performance and digestive metabolism in juvenile black carp (*Mylopharyngodon piceus*). The results demonstrated that fish subjected to sustained training at 2 and 4 BL s⁻¹ showed significantly higher critical swimming speed (U_{crit}) and maximum metabolic rate (MMR) over the control group. Fish subjected to sustained training at 1 and 2 BL s⁻¹ showed a significantly (30 and 54%) prolonged duration, 14 and 17% higher postprandial MO₂ increment (i.e. $\dot{M}O_{2peak}$), and 62 and 92% more energy expended on specific dynamic action (SDA), respectively, after consuming a similar meal over fish kept in nearly still water. These results suggest that (1) sustained exercise training at a higher speed (2 or 4 BL s⁻¹) had a positive influence on the aerobic swimming performance of juvenile *M. piceus*, which may be associated with improved aerobic metabolism; and (2) sustained exercise training at a lower speed (1 or 2 BL s⁻¹) resulted in elevated postprandial metabolic responses in juvenile *M. piceus*.

KEY WORDS: Sustained exercise training, Swimming performance, Specific dynamic action, *Mylopharyngodon piceus*

INTRODUCTION

Exercise training has been found to have complex effects on a variety of vertebrates, including humans (Hoppeler et al., 1985; Smart and Marwick, 2004), mammals (Musch et al., 1985; Evans and Rose, 1988), birds (Butler and Turner, 1988), reptiles (Thompson, 1997; Owerkovicz and Baudinette, 2008), amphibians (Pang et al., 2011; Miller and Camilliere, 1981) and fish (Davison, 1997; Li et al., 2016). Fish are an ideal research object of exercise training because of their habit of swimming against the current (Davison, 1997). According to the presence or

absence of oxygen demand, exercise training has been divided into continuously aerobic exercise training (once a day for ~18-24 h) and intermittent anaerobic exercise training (once or twice a day for ~5-10 min) for a few weeks up to an entire year (Bainbridge, 1962; Pearson et al., 1990; Liu et al., 2009; Li et al., 2013a,b). There is a wealth of evidence showing that exercise training has significant effects on swimming performance (Li et al., 2010a,b; Pang et al., 2011), feeding (Jørgensen and Jobling, 1993; Liu et al., 2009), growth rate (Brown et al., 2011; Li et al., 2013a,b), disease resistance (Castro et al., 2011), reproductive performance (Tsadik and Bart, 2007; Palstra et al., 2010a), behavioral characteristics (Totland et al., 1987; Van der Meulen et al., 2006) and gene expression (Martin and Johnston, 2005; Palstra et al., 2010b) in fish species. In the past, most of the research on exercise training in fish has been focused on cold-water salmonids, such as rainbow trout (*Oncorhynchus mykiss*) (Farrell et al., 1990; Larsen et al., 2012), Atlantic salmon (*Salmo salar*) (Totland et al., 1987; Castro et al., 2011) and chinook salmon (*Oncorhynchus tshawytscha*) (Gallaughier et al., 2001). However, there have been several recent studies on warm-water teleosts, such as qingbo (*Spinibarbus sinensis*) (Li et al., 2013a), rock carp (*Procypris rabaudi*) (Li et al., 2013b), common carp (*Cyprinus carpio*) (He et al., 2013), darkbarbel catfish (*Peltebagrus vachelli*) (Liu et al., 2009; Li et al., 2010a) and southern catfish (*Silurus meridionalis*) (Li et al., 2010b, 2016), thus increasing the knowledge of these species after exercise training.

Because swimming performance is closely related to the ability of a fish to obtain food, find a mate, and avoid predators or unfavorable conditions, swimming performance is considered a main character determining survival, and has garnered the attention of researchers for more than half a century in many species of fish (Brett, 1964; Blake, 2004; Cai et al., 2014). Swimming in fish is traditionally classified into three types: sustained, prolonged and burst-type swimming, in terms of the duration of swimming and the intensity at which the fish swims (Brett, 1964; Beamish, 1978; Plaut, 2001). Sustained exercise is powered by aerobic metabolism and does not result in muscular fatigue for long periods of time (typically >200 min) (Beamish, 1978). The critical swimming speed (U_{crit} ; i.e. the water speed at which a fish can no longer maintain its position or its maximum sustainable swimming speed) is the index used most widely by researchers to evaluate aerobic swimming performance in fish (Plaut, 2001; Lee et al., 2003a,b; He et al., 2013). A large number of studies have shown that in addition to the huge inter-species differences, U_{crit} in fish species is heavily influenced by many abiotic and biotic factors, such as temperature (Pang et al., 2011, 2013), dissolved oxygen level (Fu et al., 2011; Zhao et al., 2012), pH (Butler et al., 1992), salinity (Plaut, 2000),

¹Laboratory of Evolutionary Physiology and Behavior, Chongqing Key Laboratory of Animal Biology, Chongqing Normal University, Chongqing 400047, China. ²Key Laboratory of Freshwater Fish Reproduction and Development (Education Ministry), Key Laboratory of Aquatic Science of Chongqing, School of Life Sciences, Southwest University, Chongqing 400715, China.

*Author for correspondence (shijianfu9@hotmail.com)

 S.F., 0000-0001-7665-9037

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Received 3 January 2018; Accepted 16 January 2018

gastrointestinal fullness (Li et al., 2010a,b), nutritional status (Zhao et al., 2012), sex (Oufiero et al., 2011), productive stage (Plaut, 2002) and predator stress (Fu et al., 2015). Numerous studies have found that exercise training is a powerful stimulus for cardio-respiratory capacity and muscle hypertrophy and hyperplasia (Davison and Goldspink, 1977; Davie et al., 1986; Farrell et al., 1991; Liu et al., 2009; Fu et al., 2011). Therefore, many fish species have shown improved aerobic swimming performance after moderate exercise training (Young and Cech, 1993; Liu et al., 2009; Li et al., 2010a,b; He et al., 2013). However, this increased U_{crit} has not been found in other trained fish species (Farrell et al., 1991; Gruber and Dickson, 1997; Gallagher et al., 2001), which could be mainly due to the different species, training regimes and other environmental factors such as temperature in the training process (Davison, 1997; Pang et al., 2013).

In addition to swimming, feeding (and hence digestion) is also an important physiological function for any fish species (McCue, 2006). Specific dynamic action (SDA) is the term used to refer to the increased oxygen consumption rate ($\dot{M}O_2$) that occurs in postprandial animals, which represents the total energy expended on activities associated with the capture, handling, ingestion, digestion, absorption and assimilation of a meal, and protein synthesis and deposition associated with growth (Jobling, 1981; Beamish and Trippel, 1990; Brown and Cameron, 1991). The SDA of fish could be dependent on the species (Fu et al., 2009) and is strongly influenced by a variety of extrinsic factors, such as dietary composition (Fu et al., 2005a; Fu, 2007), meal size (Fu et al., 2005b; Wang et al., 2012), fasting (Fu et al., 2005c), temperature (Pirozzi and Booth, 2009; Pang et al., 2011) and dissolved oxygen content (Jordan and Steffensen, 2007; Fu et al., 2011). Some improvements in the cardio-respiratory system, such as the pumping performance of the heart, lamellar surface area of the gill and hemoglobin concentration of the blood were seen in trained fish, which may have some significant effects on the digestive system (Farrell et al., 1990; Gallagher et al., 2001; Gamperl and Farrell, 2004; Liu et al., 2009; Fu et al., 2011). Indeed, several previous studies found that exercise training has a profound effect on the postprandial metabolic response in fish species (Li et al., 2010a,b, 2013a,b). However, it is still debated whether exercise training could improve the postprandial metabolism in different species of fish.

In this study, we selected juvenile black carp (*Mylopharyngodon piceus*), a warm-water and benthic cyprinid fish, as the experimental animal. *M. piceus* is widely distributed in eastern Asia and is one of the four most important cultured fish species in Chinese aquaculture history (the four major Chinese carp) (Liu et al., 2004). Larvae and small juveniles feed on zooplankton and aquatic insects in the environment, where variations in natural water velocity frequently occur due to historic cycles of flood and drought, whereas the water velocity is often severely altered by dams, flood-control projects and other human activities (Chen et al., 2014). The crucial physiological

functions of all of the aquatic organisms, including fish in rivers and streams, might have corresponding effects. The objectives of this study are to test (1) whether sustained exercise training has effects on swimming performance and the postprandial metabolic response (i.e. SDA), and whether the possible effects varied with water velocity or differed between swimming and digestion; and (2) the possible underlying mechanism related to cardio-respiratory capacity and swimming efficiency. To achieve these aims, we assessed sustainable swimming performance by U_{crit} , cardio-respiratory capacity by maximum metabolic rate (MMR) and the relative sizes of the heart and gill, the swimming efficiency by the cost of transport (COT), and, finally, the postprandial $\dot{M}O_2$ responses in juvenile *M. piceus* after sustained exercise training with different velocities.

RESULTS

Effects of sustained exercise training on U_{crit} , cardio-respiratory capacity and swimming efficiency

The fish in the 2 and 4 BL s^{-1} training groups showed a significantly higher U_{crit} than those in the control and 1 BL s^{-1} groups ($P < 0.05$) (Table 1). Although neither the heart index nor the gill index showed significant differences among the experimental groups, the MMRs of both the 2 and 4 BL s^{-1} groups were significantly higher than that of the control group ($P < 0.05$), whereas the MMR of the 4 BL s^{-1} group was also significantly higher than that of the 1 BL s^{-1} group.

$\dot{M}O_2$ increased significantly with increases in swimming speed (Fig. 1). The COT significantly decreased and then reached a plateau with an increase in the swimming speed for all training and control groups (Fig. 2). Sustained exercise training and swimming speed had significant effects on the $\dot{M}O_2$ of the fish ($P < 0.05$). Sustained exercise training produced no significant effect on the COT, whereas the swimming speed had a significant effect on the COT of the fish ($P < 0.001$, Table 2).

Effects of sustained exercise training on the postprandial metabolic response

There were no significant differences in the $\dot{M}O_2$ within each time point between the ungavaged and sham-gavaged groups (Fig. 3). The postprandial $\dot{M}O_2$ of both trained and control fish increased immediately after feeding and then slowly returned to pre-fed levels (Fig. 4). There were no significant differences in the body mass, resting metabolic rate (RMR) and time to peak $\dot{M}O_2$ (PMR) among the four groups (Table 3). The SDA durations were significantly longer for the 1 and 2 BL s^{-1} training groups than those for the control and 4 BL s^{-1} training groups ($P < 0.05$). The 1 and 2 BL s^{-1} groups showed a significantly higher PMR and factorial metabolic scope over the control group ($P < 0.05$), whereas the PMR and factorial metabolic scope of the 4 BL s^{-1} group were not significantly different from those of the other three groups. The 1

Table 1. Effects of exercise training on several variables related to critical swimming performance in juvenile black carp (*M. piceus*)

	Control group	1 BL s^{-1} training group	2 BL s^{-1} training group	4 BL s^{-1} training group
Sample number (<i>n</i>)	8	8	8	8
Body mass (g)	14.94±0.95	15.20±0.65	15.36±0.76	15.68±0.84
Body length (cm)	9.16±0.20	9.31±0.15	9.36±0.19	9.26±0.15
U_{crit} (cm s^{-1})	59.39±1.91 ^b	60.96±3.01 ^b	70.98±0.67 ^a	68.00±2.34 ^a
MMR (mgO ₂ kg ⁻¹ h ⁻¹)	917.42±56.22 ^c	953.85±47.66 ^{bc}	1107.94±25.21 ^a	1060.55±52.71 ^{ab}
Heart index (%)	1.18±0.06	1.18±0.06	1.17±0.03	1.22±0.04
Gill index (%)	11.77±0.42	12.17±0.37	11.91±0.44	12.00±0.39

Data are presented as the mean±s.e.m. ^{a,b,c}Values in each row without a common lowercase letter are significantly different ($P < 0.05$).




Fig. 1. The effects of swimming speed on the oxygen consumption rate MO_2 of juvenile black carp (*M. piceus*) in the control and exercise training groups. $n=8$ for all experimental groups. Data are presented as mean \pm s.e.m. Open blue rhombuses, control group; filled pink triangles, 1 BL s^{-1} training group; filled green circles, 2 BL s^{-1} training group; filled purple squares, 4 BL s^{-1} training group.

and 2 BL s^{-1} groups showed a significantly higher energy expenditure during SDA and the SDA coefficients over the control group, whereas both variables of the 2 BL s^{-1} group were also significantly higher than those of the 4 BL s^{-1} group ($P<0.05$). However, the energy expended during SDA and the SDA coefficients of the 4 BL s^{-1} group were not significantly different from the control and 1 BL s^{-1} groups.

DISCUSSION

Effects of sustained exercise training on the swimming performance of juvenile *M. piceus*

Because sustained exercise training (aerobic training) typically involves swimming speeds that mainly utilize red muscle, it has been suggested that aerobic endurance training has a significant effect on aerobic swimming performance among fish species (Davison, 1994, 1997). In this study, 8 weeks of sustained exercise training at water velocities of 2 or 4 BL s^{-1} resulted in an 18 or 13% increase in U_{crit} for juvenile *M. piceus*, respectively. This result is similar to findings previously documented for other fish species. For example, a 30% increase in U_{crit} was found in trained striped bass




Fig. 2. The effects of swimming speed on the COT of juvenile black carp (*M. piceus*) in the control and exercise training groups. $n=8$ for all experimental groups. Data are presented as mean \pm s.e.m. Open blue rhombuses, control group; filled pink triangles, 1 BL s^{-1} training group; filled green circles, 2 BL s^{-1} training group; filled purple squares, 4 BL s^{-1} training group.

Table 2. The effect of sustained exercise training and swimming speed on MO_2 and COT in juvenile black carp (*M. piceus*)

	Training effect	Swimming speed effect	Interaction effect
MO_2	$F_{3,185}=3.017$ $P=0.031^*$	$F_{7,185}=166.704$ $P<0.001^*$	$F_{19,185}=0.320$ $P=0.997$
COT	$F_{3,185}=2.099$ $P=0.102$	$F_{7,185}=57.392$ $P<0.001^*$	$F_{19,185}=0.394$ $P=0.990$

* $P<0.05$ (based on two factors).

(*Morone saxatilis*) at 1.2 to 2.4 BL s^{-1} for 60 days (Young and Cech, 1993). In qingbo (*Spinibarbus sinensis*), 14 days of sustained training at 60% U_{crit} resulted in a 7% increase in U_{crit} (Zhao et al., 2012). Improved U_{crit} was also recorded in trained common carp (*Cyprinus carpio*) (60% U_{crit} for 28 days) (He et al., 2013) and goldfish (*Carassius auratus*) (70% U_{crit} for 48 h) (Fu et al., 2011). However, such an effect is known to be strongly dependent on the fish species, training speed and training duration (Liu et al., 2009). For example, exercise training at ~ 0.7 BL s^{-1} for 6 weeks showed no significant effect on U_{crit} in leopard shark (*Triakis semifasciata*) (Gruber and Dickson, 1997), and a U_{crit} swim test on alternate days for 4 months had no significant differences in trained chinook salmon (*Oncorhynchus tshawytscha*) compared with untrained fish (Gallaughier et al., 2001). It is worthy to note that sustained exercise training at a water velocity of 1 BL s^{-1} for 8 weeks exhibited no significant influence on U_{crit} in juvenile *M. piceus*, which suggests that the effect of sustained exercise training on U_{crit} was closely related to the training intensity in juvenile *M. piceus*.

Many studies have found that the improvement in the aerobic swimming capacity was often accompanied by increased cardio-respiratory capacity after exercise training in fish species (Farrell et al., 1991; Gallaughier et al., 2001; Liu et al., 2009; Fu et al., 2011). This was also the case in the present study, as sustained exercise training in the 2 and 4 BL s^{-1} groups resulted in a significant increase in the MMR (20 and 16%, respectively) and U_{crit} (18 or 13%, respectively) compared to the control group. Similar results were also documented in trained fish species, such as the rainbow trout (Farrell et al., 1991), striped bass (Young and Cech, 1993) and darkbarbel catfish (*Peltebagnus vachelli*) (Liu et al., 2009; Li et al., 2010a). Interestingly, heart and gill indexes in all of the trained fish had no significant difference compared to the control group in juvenile *M. piceus*, which indicated that an improved cardio-




Fig. 3. The effects of the gavage procedure on the oxygen consumption rate MO_2 in juvenile black carp (*M. piceus*). $n=8$ for the two experimental groups. Data are presented as mean \pm s.e.m. Open blue rhombuses, ungavaged group; filled pink rhombuses, sham-gavaged group.




Fig. 4. The effects of aerobic exercise training on the postprandial metabolic response in juvenile black carp (*M. piceus*). $n=11, 12, 12$ and 12 for control, 1 BL s^{-1} , 2 BL s^{-1} and 4 BL s^{-1} training groups, respectively. Data are presented as mean \pm s.e.m. Open blue rhombuses, control group; filled pink triangles, 1 BL s^{-1} training group; filled green circles, 2 BL s^{-1} training group; filled purple squares, 4 BL s^{-1} training group.

respiratory capacity was not related to eutrophy of the heart or gill. The increased lamellar surface area, enhanced cardiac enzyme activities and (or) raised hematocrit and hemoglobin levels, which have been documented in other trained fish species, might be the underlying mechanisms of improved swimming performance in juvenile *M. piceus* (Farrell et al., 1990; Gamperl and Farrell, 2004; Fu et al., 2011). Furthermore, several studies showed that fish resort to anaerobic metabolism when swimming speeds approach $\sim 60\%$ U_{crit} or more (Lee et al., 2003a,b; Zhu et al., 2010). Therefore, some researchers believe that the trained fish may recruit more anaerobic metabolism at higher swimming speeds compared to non-trained individuals, which may lead to the improvement in U_{crit} in trained fish (He et al., 2013). In this case, the $\dot{M}\text{O}_2$ of trained fish swam under certain speeds, and, hence, the apparent COT simply calculated by the $\dot{M}\text{O}_2$ of any given swimming speed should be lower than that of non-trained fish. For example, Castro et al. (2011) and Brown et al. (2011) demonstrated that trained yellowtail kingfish (*Seriola lalandi*) and Atlantic salmon at a water velocity of 0.75 to 0.80 BL s^{-1} for 6 weeks exhibited lowered costs of swimming and higher swimming efficiency. However, in our study, although sustained exercise training had a significant effect on $\dot{M}\text{O}_2$, this treatment did not produce a significant effect on COT in juvenile *M. piceus*. This result indicated that the improvement in

critical swimming performance is not related to swimming efficiency after sustained exercise training in juvenile *M. piceus*. Therefore, more data regarding the effect of exercise training on the costs of swimming in different species of fish under different training regimes are needed.

The effects of sustained exercise training on the postprandial metabolic response in juvenile *M. piceus*

The minimal maintenance metabolic rate of a postabsorptive resting ectotherm, below which the physiological function is impaired, is often defined as RMR in fish species. This physiological parameter is usually estimated from measurements of $\dot{M}\text{O}_2$ and represents the basic cost of living including ventilation, circulating body fluids, protein synthesis, and maintaining ionic gradients and osmotic work (Johnston, 1993; Norin and Malte, 2011). To date, there is no general consensus as to the effects of exercise training on the RMR of fish species. Many of the apparent discrepancies are generally caused by differences in the fish species chosen and the regime of training utilized. For instance, a higher RMR was found in trained fish, such as qingbo (sustained training at $1, 2$ and 4 BL s^{-1} for 8 weeks) (Li et al., 2013a), common carp (sustained training at $60 U_{\text{crit}}$ for 4 weeks) (He et al., 2013) and zebrafish (*Danio rerio*) (sustained training at 5 BL s^{-1} for 8 or 11 days) (Brown et al., 2011). However, in rock carp (*Procypris rabaudi*), exhaustive chasing training for 21 days had a negative effect on RMR (Li et al., 2013b). A decreased RMR in trained fish has also been documented in rainbow trout when subjected to a water velocity of 0.9 BL s^{-1} for 9 weeks compared to those reared in still water (Skov et al., 2011; Larsen et al., 2012). In this study, sustained exercise training at $1, 2$ and 4 BL s^{-1} for 8 weeks did not produce any effect on RMR. Similar findings were also demonstrated in fish species, such as southern catfish subjected to sustained training at 1 and 2 BL s^{-1} for 8 weeks (Li et al., 2016), darkbarbel catfish subjected to exhaustive chasing training for 14 days (Liu et al., 2009) and yellowtail kingfish subjected to sustained training at 0.75 BL s^{-1} for 6 weeks (Brown et al., 2011).

It is often believed that the postprandial PMR represents the maximum digestive metabolism in digesting fish and that a more rapid and higher PMR in SDA would be beneficial to the faster digestion of food and accretion of tissues (Jobling, 1981; Millidine et al., 2009). Until now, only a few studies have documented whether exercise training improved the PMR and was dependent on the species used in the different experiments. For example, cyprinid such as qingbo exhibited a higher PMR after exercise training, while another close related cyprinid, rock carp, exhibited a lower PMR after exercise training (Li et al., 2013a,b). Studies on catfish species

Table 3. The effects of aerobic exercise training on the postprandial metabolic response in juvenile black carp (*M. piceus*) measured by several variables

	Control group	1 BL s^{-1} training group	2 BL s^{-1} training group	4 BL s^{-1} training group
Sample number (n)	11	12	12	12
Body mass (g)	15.14 ± 1.22	14.89 ± 0.85	16.07 ± 0.72	16.31 ± 0.98
RMR ($\text{mgO}_2 \text{ kg}^{-1} \text{ h}^{-1}$)	147.08 ± 3.76	145.30 ± 5.59	146.67 ± 4.55	146.35 ± 5.63
Meal size (% body mass)	2.05 ± 0.01	2.02 ± 0.02	2.01 ± 0.01	2.03 ± 0.01
Energy ingested (kJ kg^{-1})	179.18 ± 1.20	176.71 ± 1.33	175.70 ± 1.18	177.73 ± 0.99
Duration (h)	10.27 ± 0.59^b	13.36 ± 0.99^a	15.86 ± 0.76^a	9.28 ± 0.53^b
Time to peak metabolic rate (h)	3.64 ± 0.36	3.79 ± 0.38	3.64 ± 0.33	3.14 ± 0.33
PMR ($\text{mgO}_2 \text{ kg}^{-1} \text{ h}^{-1}$)	251.07 ± 7.35^b	285.28 ± 6.12^a	293.99 ± 7.77^a	278.39 ± 8.55^{ab}
Factorial metabolic scope	1.71 ± 0.05^b	1.98 ± 0.07^a	2.01 ± 0.05^a	1.94 ± 0.09^{ab}
Energy expended on SDA (kJ kg^{-1})	7.95 ± 0.81^c	12.89 ± 1.39^{ab}	15.33 ± 0.92^a	9.34 ± 1.08^{bc}
SDA coefficient (%)	4.44 ± 0.46^c	7.30 ± 0.79^{ab}	8.74 ± 0.55^a	5.26 ± 0.61^{bc}

Data are presented as the mean \pm s.e.m. ^{a,b,c}Values in each row without a common lowercase letter are significantly different ($P < 0.05$).

found that PMR was inflexible when darkbarbel catfish and southern catfish underwent training (Li et al., 2010a,b, 2016). Though the precise reasons behind these different results are unknown, the change in PMR in cyprinids after exercise training may be partially due to the great flexibility of cardio-respiratory systems, which is the byproduct of natural selection on hypoxia tolerance in cyprinids during evolution (Nilsson and Renshaw, 2004; Fu et al., 2011, 2013). This is again supported by the observations that *M. piceus* of the 1 and 2 BL s⁻¹ training groups had a higher postprandial PMR compared with the control fish in the present study, which suggests that sustained exercise training had a positive effect on maximum digestive metabolism in juvenile *M. piceus*.

Metabolic scope is the difference between the MMR and RMR and it should set the limit for the magnitude of oxygen demanding processes that can be performed simultaneously, such as feeding and swimming (Clark et al., 2013). Some studies on fish species have been shown to have PMR close to MMR, which means that MMR limits the rate at which a meal can be digested (Soofiani and Hawkins, 1982; Armstrong et al., 1992; Li et al., 2010b). In the present study, the PMR (251–278 mgO₂ kg⁻¹ h⁻¹) was much lower than the MMR (917–1060 mgO₂ kg⁻¹ h⁻¹), which suggested that the costs of SDA and routine swimming are readily accommodated in the metabolic scope and juvenile *M. piceus* maintains a spare metabolic scope in an environment with unpredictable feeding opportunities (Armstrong and Schindler, 2011). Although MMR was increased by sustained exercise training at 4 BL s⁻¹, these exercise-trained juvenile *M. piceus* exhibited a similar postprandial PMR compared with untrained fish. It indicated that juvenile *M. piceus* showed no further improvement in maximum feeding metabolism under high intensity training conditions. These results suggested that more blood flow may be distributed to muscles than to digestive organs. It may potentially meet more oxygen demand of locomotive organs at a higher water velocity (~4 BL s⁻¹ and 60% U_{crit}).

Because more than 60% of SDA is directly attributed to the cost of protein synthesis and turnover and therefore to the metabolic cost of growth, it is commonly believed that SDA is closely related to the growth performance of animals (Brown and Cameron, 1991; Wieser, 1994). However, our previous studies have shown that the effects of exercise training on the energy expended on SDA might be related to differences in the fish species and types of exercise training. For example, studies on catfish species found lower SDA coefficients in exercise-trained darkbarbel catfish and southern catfish compared to untrained individuals after exhaustive chasing training for 21 days (Li et al., 2010a,b), whereas an identical training regime resulted in a decreased SDA coefficient in qingbo but a similar SDA coefficient in rock carp (Li et al., 2013b). Furthermore, it has been found that juvenile qingbo and southern catfish subjected to sustained exercise training showed a similar SDA coefficients compared to non-trained fish (Li et al., 2013a, 2016). In the present study, sustained exercise training caused significant increases in the energy expended on SDA and the SDA coefficient at water velocities of 1 and 2 BL s⁻¹ in juvenile *M. piceus*, which may be a result of the increased PMR and the extended duration of SDA. These results might suggest that the fish in 1 and 2 BL s⁻¹ training group were less efficient at digestion and required more energy to digest a standard meal. Another possibility is that the exercised fish digested the meal more completely and hence had a higher SDA coefficient. However, due to the growth rate and feed coefficient of the experimental fish not being measured in this study, we cannot determine which is true. Interestingly, exercise-trained juvenile *M. piceus* at 4 BL s⁻¹ exhibited a similar SDA coefficient compared with untrained fish,

which indicated that juvenile *M. piceus* subjected to high intensity training did not allocate more energy to their digestive system. These results also suggested that the effects of sustained exercise training on the energy expended on SDA were dependent on the intensity of training in juvenile *M. piceus*.

In conclusion, this study showed that sustained exercise training at a higher water velocity of 2 or 4 BL s⁻¹ increased the U_{crit} in *M. piceus*, which was at least partly due to the enhanced aerobic metabolism rather than improved swimming efficiency, compared to the controls. Moreover, the results regarding SDA (measured by the PMR and SDA coefficient of the fish) showed increases in postprandial metabolic responses in the fish trained at a low water velocity of 1 and 2 BL s⁻¹ rather than 4 BL s⁻¹.

MATERIALS AND METHODS

Experimental animals and acclimation

Juvenile *M. piceus* (Cypriniformes: Cyprinidae) were purchased from a local fisheries hatchery in Beibei, Chongqing, China. The fish were kept in a laboratory cement pit system (~1200 l) with recirculating water for 4 weeks before the experiment. During the acclimation period, the temperature of the dechlorinated freshwater system was maintained at 25.0±0.5°C and the oxygen content was controlled above 7 mg l⁻¹. The fish were fed to satiation twice daily (09:00 and 18:00) with commercial floating pellets (composition: 41.2±0.9% protein; 8.5±0.5% lipid; 25.7±1.2% carbohydrate and 12.3±0.4% ash). The photoperiod was 12 h light: 12 h dark with the lights turned on and off at 08:00 and 20:00, respectively.

Training regime

A self-made exercising system was used for training in the present study. The exercise system consisted of a water-processing and temperature-controlling system, a tank (190 cm×110 cm×25 cm, L×W×D), an experimental flume (140 cm×15 cm×20 cm, L×W×D), a propeller, a motor (30 w) and transducer power (see the structure in Li et al., 2016).

At the end of the acclimation period, 144 fish of similar size (11.90±0.24 g and 8.94±0.31 cm) were transferred into the exercising system for exercise training. These fish were randomly selected and divided into four groups: the control group, the 1 BL s⁻¹ training group, the 2 BL s⁻¹ training group and the 4 BL s⁻¹ training group (36 fish per group). These fish from different groups were placed in flumes of the exercising system with different water velocities. The fish in the control group swam at an average water velocity of 3 cm s⁻¹ (Li et al., 2013a, 2016). The water velocity guaranteed full water exchange and did not lead to an intense reaction in the juvenile *M. piceus*. The fish in the three training groups were forced to swim against three different water velocities [9 cm s⁻¹ (1 BL s⁻¹ exercise group), 18 cm s⁻¹ (2 BL s⁻¹ exercise group) and 36 cm s⁻¹ (4 BL s⁻¹ exercise group)] for 18 h per day at the beginning of the experiment. Continuous water velocities in the experimental flume were achieved via the motors (30 w) with a propeller. Different water velocities were produced by controlling the different voltages of transducer power. To maximize the homogeneity of water velocity along the flume, (1) the water-distributing units were installed at the head and tail of each water flume; (2) the bottom of the flume was made into a semi-circular shape; and (3) the tail of the flume was slightly higher than the head. Our pilot experiment found that a water velocity of 36 cm s⁻¹ is equivalent to ~60% of U_{crit} for juvenile *M. piceus*. To reduce physiological stress, the water velocity was gradually increased over 4 days until the desired water velocity was reached for the first round of training (Davison and Goldspink, 1978). The water velocities were adjusted every other week after the body length of the fish had been measured. The training was conducted for 8 weeks. The holding conditions and feeding regime for the experimental period were consistent with those of the acclimation period.

Measurement of variables related to U_{crit}

A Brett-type swimming tunnel respirometer (total volume 3.5 l; for details, see Li et al., 2010a and Pang et al., 2011) was used to measure the critical swimming speed (U_{crit}) of the fish in the present study. The respirometer was constructed from clear plastic polymethyl-methacrylate (PMMA). Eight fish

from each group were individually transferred into the swim tunnel and allowed to recover for 6 h after 24 h of fasting and then were swum downstream of the propeller in a swimming chamber with a cross-sectional area of 19.9 cm². The water velocity was increased by 9 cm s⁻¹ every 20 min until the fish fatigued. Fatigue was defined as the failure of the fish to move away from the rear honeycomb screen of the swimming chamber for 20 s (Lee et al., 2003a,b). The water temperature in the swimming chamber was controlled within 25.0±0.2°C using a water bath connected to a stainless steel heat exchanger. The U_{crit} was calculated for individual fish using Brett's equation (Brett, 1964):

$$U_{crit} = V + (t/T)\Delta V \quad (1)$$

where V is the highest speed at which the fish swam during the time of the experiment (cm s⁻¹), ΔV is the velocity increment (9 cm s⁻¹), T is the prescribed period of swimming per speed (20 min) and t is the time that the fish swam at the final speed (min). U_{crit} was corrected for the solid blocking effect if the cross-sectional area of the fish was more than 10% of the swimming chamber.

MO₂ as a function of swimming speed

The swimming tunnel respirometer was used to measure the $\dot{M}O_2$ as a function of swimming speed. The respirometer can be switched between an open mode and a closed mode for either replenishment of oxygen or measurement of $\dot{M}O_2$. In open mode, the respirometer was supplied with fully aerated and thermoregulated water that circulated in a reservoir tank at an approximate flow rate of 500 ml min⁻¹. In the closed mode, a small fraction of the water from the sealed respirometer was siphoned past the probe of an oximeter (HQ_{30d}, Hach Company, Loveland, CO, USA) in a cuvette thermoregulated with a water bath. The water oxygen concentration (mg l⁻¹) was recorded once every 2 min. The $\dot{M}O_2$ of an individual swimming fish was calculated from the depletion of oxygen according to the following equation (Li et al., 2010b):

$$\dot{M}O_2 = 60 \text{slope } VOL/m \quad (2)$$

where the slope (mgO₂ l⁻¹ min⁻¹) is the decrease in the water's dissolved oxygen content per minute. The slope was obtained with linear regressions between time (min) and the water's dissolved oxygen content (mgO₂ l⁻¹); only slopes with an $r^2 > 0.95$ were considered in the analysis. VOL is the total volume of the respirometer (3.5 l) minus the volume of the fish, and m is the body mass (kg) of the fish. The water oxygen content in the respirometer was never allowed to fall below 85% oxygen saturation (Claireaux et al., 2006). The maximum $\dot{M}O_2$ was used as the value for MMR (mgO₂ kg⁻¹ h⁻¹) during the U_{crit} test.

The COT (J kg⁻¹ m⁻¹) was calculated according to the following equation (Claireaux et al., 2006):

$$COT = \dot{M}O_2 \times OE/v \quad (3)$$

where $\dot{M}O_2$ (mg O₂ kg⁻¹ h⁻¹) is the oxygen consumption rate of an individual swimming fish at a given water velocity, OE is an oxycaloric equivalent of 13.54 J (mg O₂)⁻¹ and v (m h⁻¹) is the corresponding water velocity converted from cm s⁻¹ to m h⁻¹.

Heart and gill index

After the measurement of U_{crit} , the same eight fish were removed from the swimming chamber and euthanized with an overdose of MS-222 (tricaine methane sulfonate). The measurements of body mass and body length were collected to the nearest 0.1 cm and 0.1 g. The heart and gills were quickly removed with sharp scissors and cleared with 0.75% NaCl. Then, the surface water of the organs was dried by absorbent paper and weighed to the nearest 0.0001 g. The heart index and gill indexes were calculated for individual fish using the following equations (Li et al., 2016):

$$\text{Heart index} = (\text{heart mass} \times \text{body mass}^{-1}) \times 1000 \quad (4)$$

$$\text{Gill index} = (\text{gill mass} \times \text{body mass}^{-1}) \times 1000 \quad (5)$$

MO₂ of postprandial fish

The $\dot{M}O_2$ of the postprandial fish was measured using a continuous-flow respirometer (see the structure in Fu et al., 2005a). A gavage protocol (see the details in Li et al., 2013a) was performed because the fish did not eat food voluntarily in the respirometer chamber. To evaluate the effects of gavage treatment on the $\dot{M}O_2$ in juvenile *M. piceus*, 16 fish (four fish from each group, 15.35±1.15 g) were transferred into the respirometer chamber after 24 h of fasting and allowed to acclimate for another 48 h. The $\dot{M}O_2$ was measured four times in 1-h intervals before treatment. Eight fish (two fish from each group) were gently removed from the respirometer chamber and anesthetized (neutralized MS222, 50 mg l⁻¹) for ~2-3 min in a small container until they lost normal reflexes. The tip of a syringe (1 ml) without a needle was then inserted into the proximal intestine. However, no food was injected into the proximal intestine (sham-gavaged group). The fish were subsequently returned to the continuous-flow respirometer chamber. The remaining eight fish were not subjected to the procedure (ungavaged group). The $\dot{M}O_2$ was measured at 1-h intervals for 20 h.

To compare the postprandial $\dot{M}O_2$ response of the fish in the four groups, 12 fish from each group were transferred into the respirometer chamber after 24 h of fasting and allowed to acclimate for another 48 h. The $\dot{M}O_2$ was measured four times in 1-h intervals before feeding, and the means were defined as the RMR (Fu et al., 2005b; Li et al., 2010b). An identical gavage procedure was performed. Compound feed (pellet feed diluted at a ratio of 1:1.5 with water) was injected into the proximal intestine (2% body mass, which was the maximum meal size for voluntary feeding during acclimation) in a 1-min period. The fish were subsequently returned to the continuous-flow respirometer chamber. The $\dot{M}O_2$ was measured at 1-h intervals for 15 h (control and 4 BL s⁻¹ training groups) to 20 h (1 and 2 BL s⁻¹ training groups). One fish from the control group disgorged the compound feed they had been given during the experimental process, and the data from this fish were not included in subsequent analyses. The following formula was used to calculate $\dot{M}O_2$ (mg O₂ kg⁻¹ h⁻¹):

$$\dot{M}O_2 = \Delta O_2 \times F/m \quad (6)$$

where ΔO_2 is the difference in the oxygen concentration (mg O₂ l⁻¹) between the experimental chamber and the control chamber (the chamber without fish); F is the water flow rate in the experimental chamber (l h⁻¹); and m is the body mass of the fish (kg). The dissolved oxygen concentration was measured at the outlet of the chamber using an oxymeter (HQ30d, Hach Company, Loveland, CO, USA). The flow rate of water through the respirometer chamber was measured by collecting the water that was expelled from each chamber. The flow rate of each chamber was adjusted to assure 70% saturation of dissolved oxygen in the water exiting the chamber to avoid undue stress on the physiology of the fish (Blaikie and Kerr, 1996; Fu et al., 2005a). All of the experiments were conducted under constant light to minimize the effect of the circadian rhythm on fish $\dot{M}O_2$ (Fu et al., 2005a).

We quantified the following parameters for the description of SDA: (1) RMR, the mean of three $\dot{M}O_2$ values before force-feeding; (2) the peak $\dot{M}O_2$ (PMR), which is defined as the observed maximum O₂ uptake rate in the SDA process; (3) the time to peak metabolic rate, which is calculated as the time postfeeding when the $\dot{M}O_2$ was at PMR; (4) the factorial metabolic scope, which is calculated as PMR divided by RMR; (5) the duration, which is calculated as the time postfeeding when the $\dot{M}O_2$ was not significantly different from the pre-fed level; (6) the energy expended during SDA, which is calculated as the total $\dot{M}O_2$ above RMR during the duration of SDA; and (7) the SDA coefficient (%), which is the energy expended on SDA and quantified as a percentage of the energy content of the compound feed (8.75 kJ g⁻¹). The oxygen consumption was converted to energy using a conversion factor of 13.54 J (mg O₂)⁻¹.

Statistical analysis

Statistical analyses were conducted with Excel software (Microsoft Corporation, 2003) and SPSS 17.0 software (IBM, 2008). The effects of sustained exercise training and swimming speed on $\dot{M}O_2$ and COT were assessed using a two-way analysis of variance (ANOVA). The effects of sustained exercise training on the other variables were assessed using a one-

way ANOVA. ANOVA was followed by a least-significant-difference multiple-comparison test when appropriate. The effects of the gavage procedure on the $\dot{M}O_2$ within each time point between the ungavaged and sham-gavaged groups were assessed using a *t*-test. All values were calculated as the mean \pm s.e.m., and $P < 0.05$ was used as the level of statistical significance.

Acknowledgements

We thank Jianming Yuan for help with fish collection.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: S.F.; Methodology: Xiuming Li, Xiaojin Li, H.Z., J.P., S.F.; Software: H.Z., S.F.; Investigation: J.P.; Data curation: Xiaojin Li, S.F.; Writing - original draft: Xiuming Li, Y.Z., S.F.; Funding acquisition: Xiuming Li, Y.Z., S.F.

Funding

This work was supported by the National Natural Science Foundation of China [31302160, 31670418]; the China Postdoctoral Science Foundation [2014M562271]; the 111 Project [B14037] and the Basic and Frontier Research Project of Chongqing [cstc2014yykfc80001, cstc2017jcyjAX0347] of Chongqing Science and Technology Commission; the Science and Technology Research Project of the Chongqing Education Committee [KJ1400506]; and the Foundation Project of Chongqing Normal University [14XLB001].

References

- Armstrong, J. B. and Schindler, D. E. (2011). Excess digestive capacity in predators reflects a life of feast and famine. *Nature* **476**, 84-87.
- Armstrong, J. D., Priede, I. G. and Lucas, M. C. (1992). The link between respiratory capacity and changing metabolic demands during growth of northern pike, *Esox lucius* L. *J. Fish Biol.* **41**, 65-75.
- Bainbridge, R. (1962). Training, speed and stamina in trout. *J. Exp. Biol.* **39**, 537-555.
- Beamish, F. W. H. (1978). Swimming capacity. In *Fish Physiology* (ed. S. Hoar and D. J. Randall), pp. 101-138. New York: Academic Press, Inc.
- Beamish, F. W. H. and Trippel, E. A. (1990). Heat increment: a static or dynamic dimension in bioenergetic models? *Trans. Am. Fish. Soc.* **119**, 649-661.
- Blaikie, H. B. and Kerr, S. R. (1996). Effect of activity level on apparent heat increment in Atlantic cod, *Gadus morhua*. *Can. J. Fish. Aquat. Sci.* **53**, 2093-2099.
- Blake, R. W. (2004). Fish functional design and swimming performance. *J. Fish. Biol.* **65**, 1193-1222.
- Brett, J. R. (1964). The respiratory metabolism and swimming performance of young sockeye salmon. *J. Fish. Res. Board. Can.* **21**, 1183-1226.
- Brown, C. R. and Cameron, J. N. (1991). The relationship between specific dynamic action (SDA) and protein synthesis rates in the channel catfish. *Physiol. Zool.* **64**, 298-309.
- Brown, E. J., Bruce, M., Pether, S. and Herbert, N. A. (2011). Do swimming fish always grow fast? Investigating the magnitude and physiological basis of exercise-induced growth in juvenile New Zealand yellowtail kingfish, *Seriola lalandi*. *Fish Physiol. Biochem.* **37**, 327-336.
- Butler, P. J. and Turner, D. L. (1988). Effect of training on maximal oxygen uptake and aerobic capacity of locomotory muscles in tufted ducks, *Aythya fuligula*. *J. Physiol.* **401**, 347-359.
- Butler, P. J., Day, N. and Namba, K. (1992). Interactive effects of seasonal temperature and low pH on resting oxygen uptake and swimming performance of adult brown trout *Salmo trutta*. *J. Exp. Biol.* **165**, 195-212.
- Cai, L., Chen, L., Johnson, D., Gao, Y., Mandal, P., Fang, M., Tu, Z. and Huang, Y. (2014). Integrating water flow, locomotor performance and respiration of Chinese sturgeon during multiple fatigue-recovery cycles. *PLoS ONE* **9**, e94345.
- Castro, V., Grisdale-Helland, B., Helland, S. J., Kristensen, T., Jørgensen, S. M., Helgerud, J., Claireaux, G., Farrell, A. P., Krasnov, A. and Takle, H. (2011). Aerobic training stimulates growth and promotes disease resistance in Atlantic salmon (*Salmo salar*). *Comp. Biochem. Physiol. A* **160**, 278-290.
- Chen, J. M., Shen, B. Q., Pan, Q., Sun, L. H., Jiang, J. H. and Ye, J. Y. (2014). Effects of dietary protein and lipid on growth performance and body composition of advanced fingerling black carp (*Mylopharyngodon piceus*). *Acta Hydrobiol. Sin.* **38**, 699-705.
- Claireaux, G., Couturier, C. and Groison, A.-L. (2006). Effect of temperature on maximum swimming speed and cost of transport in juvenile European sea bass (*Dicentrarchus labrax*). *J. Exp. Biol.* **209**, 3420-3428.
- Clark, T. D., Sandblom, E. and Jutfelt, F. (2013). Aerobic scope measurements of fishes in an era of climate change: respirometry, relevance and recommendations. *J. Exp. Biol.* **216**, 2771-2782.
- Davie, P. S., Wells, R. M. G. and Tetens, V. (1986). Effects of sustained swimming on rainbow trout muscle structure, blood oxygen transport, and lactate dehydrogenase isozymes: evidence for increased aerobic capacity of white muscle. *J. Exp. Zool.* **237**, 159-171.
- Davison, W. (1994). Exercise training in the banded wrasse *Notolabrus fucicola* affects muscle fibre diameter but not muscle mitochondrial morphology. *N. Z. Nat. Sci.* **21**, 11-16.
- Davison, W. (1997). The effects of exercise training on teleost fish, a review of recent literature. *Comp. Biochem. Physiol. A* **117**, 67-75.
- Davison, W. and Goldspink, G. (1977). The effect of prolonged exercise on the lateral musculature of the brown trout (*Salmo trutta*). *J. Exp. Biol.* **70**, 1-12.
- Davison, W. and Goldspink, G. (1978). The effect of training on the swimming muscles of the goldfish (*Carassius auratus*). *J. Exp. Biol.* **74**, 115-122.
- Evans, D. L. and Rose, R. J. (1988). Cardiovascular and respiratory responses to submaximal exercise training in the thoroughbred horse. *Pflüg. Arch.* **411**, 316-321.
- Farrell, A. P., Johansen, J. A., Steffensen, J. F., Moyes, C. D., West, T. G. and Suarez, R. K. (1990). Effects of exercise training and coronary ablation on swimming performance, heart size, and cardiac enzymes in rainbow trout, *Oncorhynchus mykiss*. *Can. J. Zool.* **68**, 1174-1179.
- Farrell, A. P., Johansen, J. A. and Suarez, R. K. (1991). Effects of exercise-training on cardiac performance and muscle enzymes in rainbow trout, *Oncorhynchus mykiss*. *Fish Physiol. Biochem.* **9**, 303-312.
- Fu, S.-J. (2007). The specific dynamic action of southern catfish, *Silurus meridionalis* Chen fed diets containing either raw or precooked corn starch or glucose. *Fish Physiol. Biochem.* **33**, 135-141.
- Fu, S.-J., Xie, X.-J. and Cao, Z.-D. (2005a). Effect of dietary composition on specific dynamic action in southern catfish, *Silurus meridionalis*. *Aquac. Res.* **36**, 1384-1390.
- Fu, S. J., Xie, X. J. and Cao, Z. D. (2005b). Effect of meal size on postprandial metabolic response in southern catfish (*Silurus meridionalis*). *Comp. Biochem. Physiol. A* **140**, 445-451.
- Fu, S. J., Xie, X. J. and Cao, Z. D. (2005c). Effect of fasting on resting metabolic rate and postprandial metabolic response in *Silurus meridionalis*. *J. Fish Biol.* **67**, 279-285.
- Fu, S. J., Zeng, L. Q., Li, X. M., Pang, X., Cao, Z. D., Peng, J. L. and Wang, Y. X. (2009). The behavioral, digestive and metabolic characteristics in fishes with different foraging strategies. *J. Exp. Biol.* **179**, 509-517.
- Fu, S.-J., Brauner, C. J., Cao, Z.-D., Richards, J. G., Peng, J.-L., Dhillon, R. and Wang, Y.-X. (2011). The effect of acclimation to hypoxia and sustained exercise on subsequent hypoxia tolerance and swimming performance in goldfish (*Carassius auratus*). *J. Exp. Biol.* **214**, 2080-2088.
- Fu, C., Cao, Z.-D. and Fu, S.-J. (2013). The effects of caudal fin amputation on metabolic interaction between digestion and locomotion in juveniles of three cyprinid fish species with different metabolic modes. *Comp. Biochem. Physiol. A* **164**, 456-465.
- Fu, C., Fu, S.-J., Yuan, X.-Z. and Cao, Z.-D. (2015). Predator-driven intra-species variation in locomotion, metabolism and water velocity preference in pale chub (*Zacco platypus*) along a river. *J. Exp. Biol.* **218**, 255-264.
- Gallaugh, P. E., Thorarensen, H., Kiessling, A. and Farrell, A. P. (2001). Effects of high intensity exercise training on cardiovascular function, oxygen uptake, internal oxygen transport and osmotic balance in chinook salmon (*Oncorhynchus tshawytscha*) during critical speed swimming. *J. Exp. Biol.* **204**, 2861-2872.
- Gamperl, A. K. and Farrell, A. P. (2004). Cardiac plasticity in fishes: environmental influences and intraspecific differences. *J. Exp. Biol.* **207**, 2539-2550.
- Gruber, S. J. and Dickson, K. A. (1997). Effects of endurance training in the leopard shark, *Triakis semifasciata*. *Physiol. Biochem. Zool.* **70**, 481-492.
- He, W., Xia, W., Cao, Z.-D. and Fu, S.-J. (2013). The effect of prolonged exercise training on swimming performance and the underlying biochemical mechanisms in juvenile common carp (*Cyprinus carpio*). *Comp. Biochem. Physiol. A* **166**, 308-315.
- Hoppeler, H., Howald, H., Conley, K., Lindstedt, S. L., Claassen, H., Vock, P. and Weibel, E. R. (1985). Endurance training in humans: aerobic capacity and structure of skeletal muscle. *J. Appl. Physiol.* **59**, 320-327.
- Jobling, M. (1981). The influences of feeding on the metabolic rate of fishes: a short review. *J. Fish Biol.* **18**, 385-400.
- Johnston, I. A. (1993). Growth and metabolism in Antarctic fish. *Antarct. Spec. Top.* **2**, 141-150.
- Jordan, A. D. and Steffensen, J. F. (2007). Effects of ration size and hypoxia on specific dynamic action in the cod. *Physiol. Biochem. Zool.* **80**, 178-185.
- Jørgensen, E. H. and Jobling, M. (1993). Feeding in darkness eliminates density-dependent growth suppression in Arctic charr. *Aquaculture* **1**, 90-93.
- Larsen, B. K., Skov, P. V., McKenzie, D. J. and Jørgensen, A. (2012). The effects of stocking density and low level sustained exercise on the energetic efficiency of rainbow trout (*Oncorhynchus mykiss*) reared at 19°C. *Aquaculture* **324-325**, 226-233.
- Lee, C. G., Farrell, A. P., Lotto, A., MacNctt, M. J., Hinch, S. G. and Healey, M. C. (2003a). The effect of temperature on swimming performance and oxygen consumption in adult sockeye (*Oncorhynchus nerka*) and coho (*O. kisutch*) salmon stocks. *J. Exp. Biol.* **206**, 3239-3251.
- Lee, C. G., Farrell, A. P., Lotto, A., Hinch, S. G. and Healey, M. C. (2003b). Excess post-exercise oxygen consumption in adult sockeye (*Oncorhynchus nerka*) and

- coho (*O. kisutch*) salmon following critical speed swimming. *J. Exp. Biol.* **206**, 3253-3260.
- Li, X.-M., Cao, Z.-D., Peng, J.-L. and Fu, S.-J. (2010a). The effect of exercise training on the metabolic interaction between digestion and locomotion in juvenile darkbarbel catfish (*Pelteobagrus vachelli*). *Comp. Biochem. Physiol. A* **156**, 67-73.
- Li, X.-M., Cao, Z.-D. and Fu, S.-J. (2010b). The effect of exercise training on the metabolic interaction between feeding and locomotion in the juvenile southern catfish (*Silurus meridionalis* Chen). *J. Exp. Zool. A* **313A**, 557-563.
- Li, X.-M., Yu, L.-J., Wang, C., Zeng, L.-Q., Cao, Z.-D., Fu, S.-J. and Zhang, Y.-G. (2013a). The effect of aerobic exercise training on growth performance, digestive enzyme activities and postprandial metabolic response in juvenile qingbo (*Spinibarbus sinensis*). *Comp. Biochem. Physiol. A* **166**, 8-16.
- Li, X.-M., Yu, L. J., Cao, Z. D., Fu, S. J. and Zhang, Y. G. (2013b). The effects of exhaustive chasing training on the growth performance and postprandial metabolic response in juvenile qingbo (*Spinibarbus sinensis*) and rock carp (*Procypris rabaudi*). *Freshw. Fish.* **43**, 63-68.
- Li, X.-M., Liu, L., Yuan, J.-M., Xiao, Y.-Y., Fu, S.-J. and Zhang, Y.-G. (2016). The effect of aerobic exercise and starvation on growth performance and postprandial metabolic response in juvenile southern catfish (*Silurus meridionalis*). *Comp. Biochem. Physiol. A* **193**, 36-44.
- Liu, S. P., Chen, D. Q., Duan, X. B., Qiu, S. L. and Huang, M. G. (2004). Monitoring of the four famous Chinese carps resources in the middle and upper reaches of the Yangtze River. *Res. Env. Yangtze Basin* **13**, 183-186.
- Liu, Y., Cao, Z.-D., Fu, S.-J., Peng, J.-L. and Wang, Y.-X. (2009). The effect of exhaustive chasing training and detraining on swimming performance in juvenile darkbarbel catfish (*Pelteobagrus vachelli*). *J. Comp. Physiol. B* **179**, 847-855.
- Martin, C. I. and Johnston, I. A. (2005). The role of myostatin and the calcineurin-signalling pathway in regulating muscle mass in response to exercise training in the rainbow trout *Oncorhynchus mykiss* Walbaum. *J. Exp. Biol.* **208**, 2083-2090.
- McCue, M. D. (2006). Specific dynamic action: a century of investigation. *Comp. Biochem. Physiol. A* **144**, 381-394.
- Miller, K. and Camilliere, J. J. (1981). Physical training improves swimming performance of the African clawed frog *Xenopus laevis*. *Herpetologica* **37**, 1-10.
- Millidine, K. J., Armstrong, J. D. and Metcalfe, N. B. (2009). Juvenile salmon with high standard metabolic rates have higher energy costs but can process meals faster. *Proc. R. Soc. B Biol. Sci.* **276**, 2103-2108.
- Musch, T. I., Haidet, G. C., Ordway, G. A., Longhurst, J. C. and Mitchell, J. H. (1985). Dynamic exercise training in foxhounds. I. Oxygen consumption and hemodynamic responses. *J. Appl. Physiol.* **59**, 183-189.
- Nilsson, G. E. and Renshaw, G. M. C. (2004). Hypoxic survival strategies in two fishes: extreme anoxia tolerance in the North European crucian carp and natural hypoxic preconditioning in a coral-reef shark. *J. Exp. Biol.* **207**, 3131-3139.
- Norin, T. and Malte, H. (2011). Repeatability of standard metabolic rate, active metabolic rate and aerobic scope in young brown trout during a period of moderate food availability. *J. Exp. Biol.* **214**, 1668-1675.
- Ouffero, C. E., Walsh, M. R., Reznick, D. N. and Garland, T. Jr. (2011). Swimming performance trade-offs across a gradient in community composition in Trinidadian killifish (*Rivulus hartii*). *Ecology* **92**, 170-179.
- Owerkowicz, T. and Baudinette, R. V. (2008). Exercise training enhances aerobic capacity in juvenile estuarine crocodiles (*Crocodylus porosus*). *Comp. Biochem. Physiol. A* **150**, 211-216.
- Palstra, A. P., Crespo, D., Van den Thillart, G. E. and Planas, J. V. (2010a). Saving energy to fuel exercise: swimming suppresses oocyte development and downregulates ovarian transcriptomic response of rainbow trout *Oncorhynchus mykiss*. *Am. J. Physiol. Regul. Integr.* **299**, 486-499.
- Palstra, A. P., Tudorache, C., Rovira, M., Brittiijn, S. A., Burgerhout, E., van den Thillart, G. E. J. M., Spaik, H. P. and Planas, J. V. (2010b). Establishing zebrafish as a novel exercise model: swimming economy, swimming-enhanced growth and muscle growth marker gene expression. *PLoS ONE* **5**, e14483.
- Pang, X., Cao, Z.-D. and Fu, S.-J. (2011). The effects of temperature on metabolic interaction between digestion and locomotion in juveniles of three cyprinid fish (*Carassius auratus*, *Cyprinus carpio* and *Spinibarbus sinensis*). *Comp. Biochem. Physiol. A* **159**, 253-260.
- Pang, X., Yuan, X.-Z., Cao, Z.-D. and Fu, S.-J. (2013). The effects of temperature and exercise training on swimming performance in juvenile qingbo (*Spinibarbus sinensis*). *J. Comp. Physiol. B* **183**, 99-108.
- Pearson, M. P., Spriet, L. L. and Stevens, E. D. (1990). Effect of sprint training on swim performance and white muscle metabolism during exercise and recovery in rainbow trout (*Salmo gairdneri*). *J. Exp. Biol.* **149**, 45-60.
- Pirozzi, I. and Booth, M. A. (2009). The effect of temperature and body weight on the routine metabolic rate and postprandial metabolic response in mulloway, *Argyrosomus japonicus*. *Comp. Biochem. Physiol. A* **154**, 110-118.
- Plaut, I. (2000). Resting metabolic rate, critical swimming speed, and routine activity of the euryhaline cyprinodontid, *Aphanius dispar*, acclimated to a wide range of salinities. *Physiol. Biochem. Zool.* **73**, 590-596.
- Plaut, I. (2001). Critical swimming speed: its ecological relevance. *Comp. Biochem. Physiol. A* **131**, 41-50.
- Plaut, I. (2002). Does pregnancy affect swimming performance of female Mosquitofish, *Gambusia affinis*? *Funct. Ecol.* **16**, 290-295.
- Skov, P. V., Larsen, B. K., Frisk, M. and Jokumsen, A. (2011). Effects of rearing density and water current on the respiratory physiology and haematology in rainbow trout, *Oncorhynchus mykiss* at high temperature. *Aquaculture* **319**, 446-452.
- Smart, N. and Marwick, T. H. (2004). Exercise training for patients with heart failure: a systematic review of factors that improve mortality and morbidity. *Am. J. Med.* **116**, 693-706.
- Soofiani, N. M. and Hawkins, A. D. (1982). Energetic costs at different levels of feeding in juvenile cod, *Gadus morhua* L. *J. Fish Biol.* **21**, 577-592.
- Thompson, G. G. (1997). Do training and captivity affect maximal metabolic rate of *Varanus gouldii* (*Squamata: Varanidae*)? *Amphib. Reptilia* **18**, 112-116.
- Totland, G. K., Kryvi, H., Jødestøl, K., Christiansen, E. N., Tangerås, A. and Stinde, E. (1987). Growth and composition of the swimming muscle of adult Atlantic salmon (*Salmo salar* L.) during long-term sustained swimming. *Aquaculture* **66**, 299-313.
- Tsodik, G. G. and Bart, A. N. (2007). Effects of feeding, stocking density and water-flow rate on fecundity, spawning frequency and egg quality of Nile tilapia, *Oreochromis niloticus* (L.). *Aquaculture* **272**, 380-388.
- Van der Meulen, T., Schipper, H., van den Boogaart, J. G. M., Huising, M. O., Kranenbarg, S. and Van Leeuwen, J. L. (2006). Endurance exercise differentially stimulates heart and axial muscle development in zebrafish (*Danio rerio*). *Am. J. Physiol. Regul. Integr.* **291**, 1040-1048.
- Wang, Q., Wang, W., Huang, Q., Zhang, Y. and Luo, Y. (2012). Effect of meal size on the specific dynamic action of the juvenile snakehead (*Channa argus*). *Comp. Biochem. Physiol. A* **161**, 401-405.
- Wieser, W. (1994). Cost of growth in cells and organisms: general rules and comparative aspects. *Biol. Rev. Camb. Philos. Soc.* **69**, 1-33.
- Young, P. S. and Cech, J. J., Jr. (1993). Improved growth, swimming performance, and muscular development in exercised-conditioned young-of-the-year striped bass (*Morone saxatilis*). *Can. J. Fish. Aquat. Sci.* **50**, 703-707.
- Zhao, W.-W., Pang, X., Peng, J.-L., Cao, Z.-D. and Fu, S.-J. (2012). The effects of hypoxia acclimation, exercise training and fasting on swimming performance in juvenile qingbo (*Spinibarbus sinensis*). *Fish Physiol. Biochem.* **38**, 1367-1377.
- Zhu, Y.-P., Cao, Z.-D. and Fu, S.-J. (2010). Aerobic and anaerobic metabolism in response to different swimming speed of juvenile darkbarbel catfish (*pelteobagrus vachelli richardson*). *Acta Hydrobiol. Sin.* **34**, 905-912.