

pH controls spermatozoa motility in the Pacific oyster (*Crassostrea gigas*)

Myrina Boulais^{1,*}, Marc Suquet², Eve Julie Arsenault-Pernet², Florent Malo², Isabelle Queau², Patricia Pignet³, Dominique Ratiskol², Jacqueline Le Grand², Matthias Huber², Jacky Cosson⁴

¹CNRS, UMR 6539 Lemar (UBO-CNRS-IRD-Ifremer), IUEM, Plouzané, France, ²Ifremer, UMR 6539 Lemar (UBO-CNRS-IRD-Ifremer), Site expérimental d'Argenton, Landunvez, France, ³Ifremer, LMEE, Centre de Bretagne, Plouzané, France, ⁴University of South Bohemia in Ceske Budejovice, Faculty of Fisheries and Protection of Waters, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Vodnany, Czech Republic.

*Author for correspondence (myrina.boulais@gmail.com; myrina.boulais@univ-brest.fr)

KEY WORDS: Seminal fluid, Spermatozoa, Motility, pH, Salinity, Ions

SUMMARY STATEMENT: Oyster sperm motility is controlled by pH and is not triggered in an acidic environment.

ABSTRACT

Investigating the roles of chemical factors stimulating and inhibiting sperm motility is required to understand the mechanisms of spermatozoa movement. In this study, we described the composition of the seminal fluid (osmotic pressure, pH, and ions) and investigated the roles of these factors and salinity in initiating spermatozoa movement in the Pacific oyster. The acidic pH of the gonad (5.82 ± 0.22) maintained sperm in the quiescent stage and initiation of flagellar movement was triggered by a sudden increase of spermatozoa external pH (pHe) when released in seawater (SW). At pH 6.4, percentage of motile spermatozoa was three times higher when they were activated in SW containing 30 mM NH_4Cl , which alkalinizes internal pH (pHi) of spermatozoa, compared to NH_4Cl -free SW, revealing the role of pHi in triggering sperm movement. Percentage of motile spermatozoa activated in Na^+ -free artificial seawater (ASW) was highly reduced compared to ASW, suggesting that change of pHi triggering sperm motility was mediated by a Na^+/H^+ exchanger. Motility and swimming speed were highest in salinities between 33.8 and 42.7‰ (within a range of 0 to 50 ‰), and pH values above 7.5 (within a range of 4.5 to 9.5).

INTRODUCTION

Movement of spermatozoa is required for external fertilization, allowing male gametes to reach and penetrate oocytes (Billard, 1983). In invertebrates and fish, sperm is non-motile in the gonad (Cosson et al., 2008; Demoy-Schneider et al., 2014). Sperm quiescence maintains energy stores of spermatozoa and prevents alterations to their morphology (Morisawa, 1985). Initiation of sperm motility takes place after sperm release into the aquatic environment and is triggered by physical or chemical changes occurring between the gonad and the aquatic environment (Alavi and Cosson, 2005, 2006). Consequently, identifying factors inhibiting and stimulating spermatozoa movement requires the description of the chemical constituents of seminal fluid and study of their roles in controlling sperm motility.

Factors activating sperm motility vary among species. In fish, the change in osmotic pressure, pH or ion composition induces sperm movement (Morisawa and Suzuki, 1980; Alavi and Cosson, 2005, 2006). In marine invertebrates, sperm activation factors have been mainly investigated in sea urchins. Rothschild (1928) first suggested that sperm motility initiation of the sea urchin, *Echinus esculentus*, depends on the “dilution effect” of spermatozoa at spawning. Later, Mohri and Yasumasu (1963) listed several factors suggested

to maintain sperm quiescence in the seminal fluid, including concentrations of CO₂, O₂ and ions (K⁺, Cu²⁺ or Zn²⁺), the presence of androgamone I, and pH. The crucial role of pH in triggering sperm motility was later confirmed in the sea urchin *Strongylocentrotus purpuratus*. In this species, seminal fluid pH and sperm intracellular pH (pHi) are more acidic than seawater (SW) (Schackmann et al. 1981; Christen et al., 1986). Finally, artificially increasing sperm pHi by adding NH₄Cl to the activating medium also triggers sperm motility (Christen et al., 1982).

In marine bivalve molluscs, knowledge of sperm biology is limited because their high fecundity compensates for the high inter-breeder variability of gamete quality, allowing mass production of marine bivalves regardless of gamete biology (Robert and Gerard, 1999). However, understanding the factors inhibiting and triggering the initiation of sperm motility is needed to support the current development of aquaculture that is oriented towards controlled crosses, selective breeding, and gamete cryobanking (Piferrer et al., 2009). As in sea urchins, acidic external pH (pHe) inhibits sperm movement in the Japanese pearl oyster (*Pinctada fucata martensii*; Ohta et al., 2007), black lip pearl oyster (*Pinctada margaritifera*; Demoy-Schneider et al., 2012, 2014), king scallop (*Pecten maximus*; Faure et al., 1994), and Pacific oyster (*Crassostrea gigas*; Dong et al., 2002; Alavi et al., 2014). A complex interaction between environmental pH and temperature modified sperm movement characteristics in the mussel, *Mytilus galloprovincialis* (Eads et al., 2016). Recently, it was shown that the concentration of some gonadal ions (i.e. K⁺, Ca²⁺, and Na⁺) are involved in controlling sperm motility in the Pacific oyster (Alavi et al., 2014). However, the roles of pHi, other ions of the seminal fluid such as Mg²⁺, the osmotic pressure, and salinity in triggering motility initiation have never been investigated in Pacific oyster spermatozoa. Furthermore, control of swimming speed, a key feature of sperm motility, by chemical factors stimulating and inhibiting motility has never been studied in bivalve species.

In this study, we described the composition of the seminal fluid (pH, ions, and osmotic pressure) and investigated the roles of these factors and salinity in initiating spermatozoa movement (percentage of motility and swimming speed) in the Pacific oyster, a model species in invertebrate reproduction.

RESULTS

Composition of the seminal fluid

Osmotic pressure, pH, and concentrations of ions of Pacific oyster seminal fluid are reported in Table 2. Seminal fluid pH of the Pacific oyster was acidic. Two ions, Cl^- and Na^+ , were predominant in the seminal fluid.

Effects of salinity, pH, and ions in initiating spermatozoa movement

Percentage of motile spermatozoa significantly increased with salinity both at 1 min after activation ($F_{1,9} = 15.88$, $P < 0.001$) and 5 min after activation ($F_{1,9} = 12.11$, $P < 0.001$) (Fig. 1A). At 5 min post activation, the highest values were recorded for salinity ranging from 19.3 to 42.7‰. Velocity of the Average Path (VAP) was not significantly different among salinities for spermatozoa activated for 1 min. However, VAP significantly increased ($F_{1,7} = 4.06$, $P < 0.01$) with salinity for spermatozoa activated for 5 min, maximum VAP values being recorded at 33.8 and 42.7‰ (Fig. 1B). Overall, there was a trend for highest motility and VAP in salinities between 33.8 and 42.7‰.

Oyster sperm was non-motile in pH values ranging from 4.5 to 5.5 (Fig. 2A). Low percentages of motile spermatozoa ($< 10\%$) were recorded at pH 6.0, 6.5, and 7.0. A significant increase of motile percentage was found at pH values of 7.5 and above, both at 1 min after activation ($F_{1,10} = 20.71$, $P < 0.001$) and 5 min after activation ($F_{1,10} = 44.94$, $P < 0.001$). For spermatozoa activated for 1 min, percentage of motile cells significantly increased between pH 8.0 and 8.5 which was not the case for spermatozoa activated for 5 min. Sperm VAP significantly increased in a pH-dependent manner from pH 6.5 for both sperm movement durations ($F_{1,10} = 11.26$ at 1 min and $F_{1,10} = 24.31$ at 5 min, $P < 0.001$ for both durations) (Fig. 2B).

Regarding the effect of increasing spermatozoa pH_i adding NH_4Cl to the activating media, percentage of motile spermatozoa at 1 min after activation was significantly lower at pH 6.4 compared to pH 8.1 when NH_4Cl was not added ($F_{1,5} = 18.71$, $P < 0.001$) (Fig. 3A). NH_4Cl (30 mM) significantly increased the percentage of motile spermatozoa activated in pH 6.4 for 1 min. Percentage of motility after 5 min of sperm activation was not significantly changed by NH_4Cl compared to controls (i.e. NH_4Cl -free SW for each corresponding pH). VAP was similar for both NH_4Cl concentrations (Fig. 3B).

VAP and percentage of motile spermatozoa were not significantly different between SW and ASW regardless of the activation time (Fig. S1). Consequently, ASW was considered as the single control for the following results. At both times after activation, a significantly lower percentage of motility ($F_{1,5} = 4.37$, $P < 0.01$ at 1 min and $F_{1,5} = 44.85$, $P < 0.001$ at 5 min after activation) was recorded for Na⁺-free media compared to ASW (Fig. 4A). Conversely, a lack of Ca²⁺ significantly increased percentage of motile spermatozoa activated for 5 min, but not for 1 min, compared to ASW. A lack of K⁺ or Mg²⁺ in activating medium did not significantly change percentage of motile spermatozoa. Sperm VAP was not significantly affected by a lack of Ca²⁺, K⁺ or Mg²⁺ in ASW (Fig. 4B, VAP in Na⁺-free ASW was not measured because of low number of motile spermatozoa). No movement of spermatozoa was observed in ion-free ASW.

DISCUSSION

The present study investigated the roles of pH, ions, and salinity in initiating Pacific oyster spermatozoa movement.

Motility of spermatozoa was triggered by a sudden alkalinization of sperm environment when released in seawater. Values of pH of 7.5 and above were optimal to initiate movement of Pacific oyster spermatozoa. This value is lower than the one suggested by Alavi et al. (2014; pHe > 8) in the same species. This is probably explained by the use of pH-indicator strips by these authors, methodology which is not as accurate as a pH meter. Although this 0.5 difference in optimal pH value between our study and the one of Alavi et al. (2014) appears small, it represents very different acidic conditions due to the logarithmic nature of the pH scale. Inhibition of sperm motility in neutral and acidic pH conditions (≤ 7.0) demonstrated that the low pH of the gonadal environment (pH 5.82) maintains sperm in the quiescent stage. The inhibitory role of acidic pH in sperm motility was also suggested in the black-lip pearl oyster, characterized by an optimal range of pH values for sperm motility ranging from 8.5 to 10.5 (Demoy-Schneider et al., 2012). In the king scallop, pHe also regulates sperm motility initiation (Faure et al., 1994).

Since pHe mediates pHi-alkalinization in spermatozoa (Christen et al., 1982), we investigated the role of pHi changes in triggering motility initiation in Pacific oyster spermatozoa. A 3-fold higher percentage of motile spermatozoa was observed when pHi was alkalinized using NH₄Cl compared to NH₄Cl-free SW. Initiation of sperm motility by alkalinization of pHi using NH₄Cl or NH₃ has been reported in the Manila clam (Alavi et al.,

2014), sea urchin species (Christen et al., 1982, 1983; Lee et al., 1983), and starfish *Asterina pectinifera* (Nakajima et al., 2005). Initiation of sperm motility by pHi alkalization is related to functionality of flagellar dynein ATPases, the enzymes sustaining axonemal movement. It was demonstrated that optimal pH values for dynein ATPase reactions ranged from 7.4 to 8.6, while their activity is inhibited at pH 7.2 (Gibbons and Gibbons, 1972; Christen et al., 1982).

Osmotic pressure of the seminal fluid was similar to the one of seawater (~ 1000 mOsm kg⁻¹ for salinity of 35 ‰; Lovett et al., 2006), demonstrating that initiation of sperm movement is osmolarity-independent in the Pacific oyster. The roles of seminal fluid ions in controlling sperm movement were explored in our work. Similarly to our results, Alavi et al. (2014) recorded no movement in ion-free ASW, demonstrating that the presence of ions is required to trigger sperm movement in this species.

Movement of Pacific oyster spermatozoa was dependent on the ionic composition of the activating medium. Sperm activation in Na⁺-free ASW decreased the percentage of motility down to 1% of the value observed in the control, showing that Na⁺ plays a major role in Pacific oyster spermatozoa movement. Dilution of spermatozoa in Na⁺-free ASW also inhibited sperm motility in the king scallop (Faure et al., 1994). In sea urchins, sperm movement is triggered by a Na⁺-dependent pHi rise suggested to be mediated by a Na⁺/H⁺ exchanger (Gatti and Christen, 1985; Darszon et al., 1999). Consequently, pHi but not Na⁺ is the primary factor controlling sperm motility in sea urchins. This statement is sustained by the initiation of spermatozoa movement following pHi alkalization using NH₄Cl (Christen et al., 1983; Lee et al., 1983). Based on the present results, a similar mechanism is suggested in Pacific oyster spermatozoa and would be of interest to address in the future.

Our results showed a higher percentage of motile spermatozoa in Ca²⁺-free ASW compared to ASW at 5 min after activation. It was previously reported that dilution of sperm in a medium lacking Ca²⁺ increased motility of Eastern oyster and Pacific oyster spermatozoa (Paniagua-Chavez et al., 1998; Dong et al., 2002). On the other hand, a lower percentage of motile spermatozoa was observed in Ca²⁺-free ASW compared to ASW in the Pacific oyster (Alavi et al., 2014). However, these authors highlighted that intracellular Ca²⁺ concentration was not the primary factor controlling motility because the addition of Ca²⁺ ionophore to sucrose- or to Na⁺-activating medium containing 10 mM Ca²⁺ did not trigger sperm motility. Plasma membrane Na⁺/Ca²⁺ exchangers are one way in which cells regulate Ca²⁺ (Blaustein and Lederer, 1999). We suggest that the higher percentage of motile spermatozoa in Ca²⁺-free ASW compared to ASW, observed in our study, is associated with changes in internal Na⁺

concentration. Investigating the roles of $\text{Na}^+/\text{Ca}^{2+}$ exchangers in controlling sperm motility initiation would be of interest for future studies.

In the present study, a lack of K^+ in the activating medium did not significantly change sperm movement characteristics of the Pacific oyster, validating that K^+ influx is not required to trigger sperm motility (Alavi et al., 2014). These authors suggested that K^+ efflux, however, is essential for initiation of sperm motility and that high concentration of K^+ in the gonad maintains sperm in the quiescent state in this species. In sea urchins, in the presence of sodium, addition of K^+ induces an internal alkalization, accompanied by an influx of potassium (Gatti and Christen, 1985).

Some differences in pH and ionic composition of seminal fluid are observed among marine invertebrates (Table 3). Spermatozoa activation is an evolutionary conserved mechanism which involves a Na^+ -dependent alkalization of pH_i in marine invertebrates. In marine fish, low osmotic pressure (306 to 400 mosmol L^{-1}) and concentration of K^+ (2.0 to 5.3 mmol L^{-1}), and high pH value (7.3 to 7.8) of the seminal fluid (reviewed in Suquet et al., 1994) reflect a different mechanism of spermatozoa activation with respect to marine invertebrates. In marine fish, pH is not involved in triggering sperm motility, but a rise of the osmotic pressure during sperm release initiates spermatozoa movement (Morisawa and Suzuki, 1980; Cosson et al., 2008).

Our results revealed that spermatozoa are equally motile in a wide range of salinities (19.3 to 42.7‰). The limited role of this environmental factor in controlling sperm motility is in agreement with optimal salinity requirements of gametogenesis and early life stage development of the Pacific oyster, ranging from 18 to 30‰ (Phelps and Warner, 1990; Helm, 2005). It was suggested that tolerance of Pacific oyster to low salinities could be sustained by molecular mechanisms of osmoadaptation (Zhao et al., 2012).

Current models predicting the degree of ocean acidification call for a drop in pH by 0.3 to 0.5 by 2100, and possibly by 0.8 to 1.4 by 2300 depending on the CO_2 emission scenario (Caldeira and Wickett, 2005; IPCC, 2013). In the present study, decreasing SW pH from 8.0 to 7.5, as predicted by the end of this century, affected neither percentage of motile spermatozoa nor swimming speed, suggesting that near-future levels of ocean acidification are not a major threat to spermatozoa motility in the Pacific oyster. Species-specific responses are found in studies investigating the effect of low pH on spermatozoa movement of marine invertebrates. In the mussel, *Mytilus galloprovincialis*, decreasing seawater pH from 8.0 to 7.6 induced a 25% reduction of percentage of motile spermatozoa and their sperm swimming speed (Vihtakari et al., 2013). In the same species, Eads et al. (2016) found that the effect of

pH on spermatozoa movement was only noticeable after 2h of exposure. In the sea urchin, *Heliocidaris erythrogramma*, seawater acidification Δ pH of 0.3 and 0.5 units resulted in 7 and 9% reduction of percentage of motile spermatozoa, respectively, without affecting sperm swimming speed (Schlegel et al., 2012). For higher levels of ocean acidification such as predicted after 2100 ($<$ pH 7.5), the movement characteristics of Pacific oyster spermatozoa will be adversely impacted, which may have profound negative implications for reproduction and subsequent recruitment in this species. Overall, oyster reproduction seems to be resilient to moderate environmental acidification. In the present study, acidification of sperm environment was obtained by manipulating pH by acid while best practice methodologies manipulate pH either by aerating or suturing seawater with CO₂ (Ross et al. 2011). Investigating the effects of ocean acidification on Pacific oyster sperm movement must be further studied, taking into account acidification methodology.

In conclusion, acidic pH of the seminal fluid compared to seawater maintains sperm in the quiescent state in the gonad of the Pacific oyster. Sperm movement is suggested to be triggered by a sudden Na⁺ dependent alkalization of pHi as a consequence of sperm release in seawater. Other cations studied in the present study on sperm motility initiation have only minor effects. The use of sperm swimming speed as a proxy of environmental conditions strengthens previous observations carried out only using the percentage of motile spermatozoa. Finally, the present observations highlight the resilience Pacific oyster sperm motility exhibits in response to ocean acidification projected for near-future.

MATERIALS AND METHODS

Animals and sperm collection

Three-year old Pacific oysters (mean weight \pm s.e.m.: 136 \pm 11 g) were collected from aquaculture stock in the Bay of Brest (Finistère, France) during their natural spawning period. They were transferred to the Argenton experimental hatchery and maintained in conditioning conditions in a 600-L tank (natural SW at 19°C, oysters fed *ad libitum* with a mixture of two microalgae, *Isochrysis galbana* and *Chaetoceros gracilis*, equal to 8% dry weight algae/dry weight oyster per day and per oyster) until their use (Fabioux et al., 2005). Since our study aimed at studying spermatozoa biological characteristics of ripe oysters, only stage 3 males were selected (Steele and Mulcahy, 1999). For each oyster, sperm was collected by sampling (700 μ L) in the proximal part of the gonad using a micropipette, and each sperm sample was stored on ice in a 2-mL plastic vial (Eppendorf Quality™) until use.

Composition of the seminal fluid

To determine pH, osmotic pressure and concentrations of ions of the seminal fluid ($n = 22$ oysters, except $n = 3$ oysters for osmotic pressure), a 500- μL subsample of sperm was centrifuged (Sigma 3-16PK) at 4,000 g for 1 hour at 4°C (1.5-mL plastic vial Eppendorf Quality™) for each individual sperm sample. Supernatant (i.e. seminal fluid) pH was immediately assessed using a pH meter (IQ150, IQ Scientific instrument; total pH scale). The supernatant was then collected in a 500- μL plastic vial (Eppendorf Quality™) and stored at –28°C until later analyses of osmotic pressure and concentrations of ions. Seminal fluid osmotic pressure was measured on a 100- μL subsample, using an automatic osmometer (Type 15, Löser). Concentrations of ions were assessed using ion chromatography (50 μL subsample, ICS 900, Thermo Scientific) fitted with a Ionpac CS 12A column and a suppressor (CSRS-ultra II) unit in combination with a DS5 conductivity cell (Thermo Scientific). Components were separated using a methanesulfonic acid eluent (18 mM), with a flow of 1 mL min^{-1} (Lazar et al., 2011).

Effects of salinity, pH, and ions in initiating spermatozoa movement

To study the effects of salinity, pH and ions on triggering movement, 5 μL of sperm was diluted in 500 μL of different activating media described below (500- μL plastic vial, Eppendorf Quality™) for each oyster ($n = 5$ oysters). Unless stated, activating media consisted of SW filtered at 1 μm , 34.5‰ salinity, 19°C, supplemented by 0.3% pluronic acid, 15 mM Tris, and final pH adjusted to 8.1 by addition of diluted HCl (all chemicals used in our experiments were purchased from Sigma Aldrich®).

The effects of salinity on sperm movement activation were explored across a range of salinities, from 0.4 to 50‰. Adjustment of salinity was conducted by dilution of activating media with distilled water or by addition of NaCl solution for salinity values higher than 35‰. The effects of pH on sperm movement initiation was tested after adjusting pH of activating media to values ranging from 4.5 to 9.5. Buffers used to adjust pH values were 15 mM acetate/acetic acid and 15 mM Tris/HCl for pH values between 4.5 and 6.0 and between 6.5 and 9.5, respectively. The effect of spermatozoa pH_i alkalization on their movement characteristics was tested at two different pH (6.4 and 8.1) adding NH_4Cl (30 mM final) as this chemical alkalizes pH_i (Christen et al., 1983).

Finally, to explore the effects of seawater ions on triggering sperm motility, different activating media were prepared, including artificial seawater (ASW) and Na⁺-, Ca²⁺-, K⁺-, Mg²⁺-, and ion-free ASW (Table 1). Ethylene glycol-bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA) was used in the ion free ASW medium because sugar may be contaminated by trace concentration of Ca²⁺. EGTA and Ethylenediaminetetraacetic acid (EDTA) had no toxic effects on sperm motility at concentration of 1 mM as these chemicals did not decrease sperm movement characteristics in Ca²⁺- and Mg²⁺-free ASW.

General procedure for assessment of sperm motility

After sperm dilution in activating medium, a 12 μL sample was transferred to a Fast read 102 cell (Biosigma). Sperm movement was recorded at the top of the drop, at 1 and 5 min after dilution in activating media, maximum values of sperm movement being previously observed after 5 min of activation (Suquet et al., 2016). Sperm samples were observed using a phase contrast microscope (Olympus BX51, x20 magnification) connected to a video camera (Qicam Fast 1934, 60 frames s⁻¹ camera). Percentage of motile spermatozoa and their VAP were assessed (n > 30 spermatozoa for each assay) using a computer-assisted sperm analyser (CASA) plugin developed for the Image J software (Wilson-Leedy and Ingermann, 2007) and calibrated to Pacific oyster according to Boulais et al. (2015).

Statistical analysis

Data were presented as mean ± s.e.m. Percentage data were arcsin square-root transformed. Normality (Shapiro-Wilk test) and homogeneity of variances (Levene's test) were verified. Data were compared using one way ANOVA. When significant differences were observed, a Fisher a posteriori test was used. Experimental results were considered significant at *P* < 0.05. Statistical analyses were performed using Statistica 6 software.

Acknowledgments

The authors want to thank the University of South Bohemia in Ceske Budejovice and the CENAKVA (South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, CZ.1.05/2.1.00/01.0024) for funding M Suquet visit. We are particularly grateful to S Pouvreau (Ifremer) for discussion on spawning behaviour of Pacific oyster in the wild and A Severe (Ifremer) for chemical advices on medium composition.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: M.B., M.S., J.C.; Methodology: E.J.A.P., F.M., I.Q., P.P., D.R., J.L.G., M.H.; Analysis: M.S., E.J.A.P., F.M., I.Q., P.P.; Writing and review: M.B., M.S., J.C.; Funding acquisition: M.S.

Funding

This work was supported by the national project CRB-anim (ANR-11-INBS-003) and the European project Reproseed (FP7-KBBE-2009-3).

References

- Alavi, S.M.H. and Cosson, J.** (2005). Sperm motility in fishes: (I) Effects of temperature and pH. *Cell Biol. Int.* **29**, 101-110. (doi: 10.1016/j.cellbi.2004.11.021)
- Alavi, S.M.H. and Cosson, J.** (2006). Sperm motility in fishes: (II) Effects of ions and osmolality. *Cell Biol. Int.* **30**, 1-14. (doi: 10.1016/j.cellbi.2005.06.004)
- Alavi, S.M.H., Matsumura, N., Shiba, K., Itoh, N., Takahashi, K.G., Inaba, K. and Osada, M.** (2014). Roles of extracellular ions and pH in 5-HT induced sperm motility in marine bivalve. *Reprod.* **147**, 331-345. (doi: 10.1530/REP-13-0418)
- Billard, R.** (1983). Ultrastructure of trout spermatozoa: changes after dilution and deep freezing. *Cell Tissue Res.* **228**, 205-218. (doi: 10.1007/BF00204873)
- Blaustein, M.P. and Lederer, W.J.** (1999). Sodium/Calcium Exchange: Its Physiological Implications. *Physiol Rev.* **79**, 763–854.
- Boulais, M., Soudant, P., Le Goïc, N., Quéré, C., Boudry, P. and Suquet, M.** (2015). Involvement of mitochondrial activity and OXPHOS in ATP synthesis during the motility phase of spermatozoa in the Pacific oyster, *Crassostrea gigas*. *Biol. Reprod.* **93**, 1-7. (doi: 10.1095/biolreprod.115.128538)
- Caldeira, K. and Wickett, M.E.** (2005). Ocean model predictions of chemistry change from carbon dioxide emissions to the atmosphere and ocean. *J. Geophys. Res.* **110**, C09S04. (doi: 10.1029/2004JC002671)
- Christen, R., Schackmann, R.W. and Shapiro, B.M.** (1982). Elevation of the intracellular pH activates respiration and motility of sperm of the sea urchin, *Strongylocentrotus purpuratus*. *J. Biol. Chem.* **257**, 14881-14890.
- Christen, R., Schackmann, R.W. and Shapiro, B.M.** (1983). Metabolism of sea urchin sperm. Interrelationships between intracellular pH, ATPase activity and mitochondrial respiration. *J. Biol. Chem.* **258**, 5392-5399.
- Christen, R., Schackmann, R.W. and Shapiro, B.M.** (1986). Ionic regulation of sea urchin sperm motility, metabolism and fertilizing capacity. *J. Physiol.* **379**, 347-365.
- Cosson, J., Groison, A.L., Suquet, M., Fauvel, C., Dreanno, C. and Billard, R.** (2008). Studying sperm motility in marine fish: an overview on the state of the art. *J. Appl. Ichthyol.* **24**, 460-486. (doi: 10.1111/j.1439-0426.2008.01151.x)
- Darszon, A., Labarca, P., Nishigaki, T. and Espinoza, F.** (1999). Calcium changes in the development, maturation and function of spermatozoa. *Physiol. Rev.* **79**, 481-510.

- Demoy-Schneider, M., Levêque, A., Schmitt, N., Le Pennec, M. and Cosson, J.** (2012) Motility activation and metabolism characteristics of spermatozoa of the black-lip-pearl oyster *Pinctada margaritifera* var: *cumingii* (Jameson 1901). *Theriogenol.* **77**, 53-64. (doi: 10.1016/j.theriogenology.2011.07.014)
- Demoy-Schneider, M., Schmitt, N., Suquet, M., Labbé, C., Boulais, M., Prokopchuk, G. and Cosson, J.** (2014). Biological characteristics of sperm in two oyster species: the Pacific oyster, *Crassostrea gigas*, and the black lip pearl oyster, *Pinctada margaritifera*. In *Spermatozoa: biology, motility and function and chromosomal abnormalities* (ed. B.T. Erickson), pp. 15-74. New York: Nova Biomedical.
- Dong, Q., Eudeline, B., Allen Jr., S.K.A. and Tiersch, T.** (2002). Factors affecting sperm motility of tetraploid Pacific oysters. *J. Shellfish Res.* **21**, 719-723.
- Eads, A.R., Kennington, W.J. and Evans, J.P.** (2016). Interactive effects of ocean warming and acidification on sperm motility and fertilization in the mussel *Mytilus galloprovincialis*. *Mar. Ecol. Progr. Ser.* **562**, 101-111.
- Fabioux, C., Huvet, A., Le Souchu, P., Le Pennec, M. and Pouvreau, S.** (2005). Temperature and photoperiod drive *Crassostrea gigas* reproductive internal clock. *Aquaculture* **250**, 458-470. (doi: 10.1016/j.aquaculture.2005.02.038)
- Faure, C., Devauchelle, N. and Girard, J.P.** (1994). Ionic factors affecting motility, respiration and fertilization rate of the sperm of the bivalve *Pecten maximus* (L.). *J. Comp. Physiol.* **164B**, 444-450. (doi: 10.1007/BF00714581)
- Gatti, J.L. and Christen, R.** (1985). Regulation of internal pH of sea urchin sperm. *J. Biol. Chem.* **260**, 7599-7602.
- Gibbons, B.H. and Gibbons, I.R.** (1972). Flagellar movement and adenosine triphosphatase activity in sea urchin sperm extracted with Triton X-100. *J. Cell Biol.* **54**, 75-97. (doi: 10.1083/jcb.54.1.75)
- Helm, M.** (2005). Cultured aquatic species information program - *Crassostrea gigas*. Cultured aquatic species fact sheets. FAO Inland Water Resources and Aquaculture Service. (http://www.fao.org/fishery/culturedspecies/Crassostrea_gigas/en#tcNA0064)
- Intergovernmental Panel on Climate Change (IPCC)** (2013). Climate change for 2013. The physical science basis (2013). Cambridge, U.K.: Cambridge University (<http://www.climatechange2013.org/>)

- Johnson, C.H., Clapper, D.L., Winkler, M.W., Lee, H.C. and Epel, D.** (1983). A volatile inhibitor immobilizes sea urchin sperm in semen by depressing the intracellular pH. *Dev. Biol.* **98**, 493-501.
- Lazar, C.S., Parkes, R.J., Cragg, B.A., L'Haridon, S. and Toffin, L.** (2011). Methanogenic diversity and activity in hypersaline sediments of the centre of the Napoli mud volcano, Eastern Mediterranean Sea. *Environ. Microbiol.* **13**, 2078-2091. (doi: 10.1111/j.1462-2920.2011.02425.x)
- Lee, H.C., Johnson, C. and Epel, D.** (1983). Changes in internal pH associated with initiation of motility and acrosome reaction of sea urchin sperm. *Dev. Biol.* **95**, 31-45. (doi: 10.1016/0012-1606(83)90004-0)
- Lovett, D.L., Tanner, C.A., Glomski, K., Ricart, T.M. and Borst, D.W.** (2006). The effect of seawater composition and osmolality on hemolymph levels of methyl farnesoate in the green crab *Carcinus maenas*. *Comp. Biochem. Physiol., Part A* **143**, 67-77. (doi: 10.1016/j.cbpa.2005.10.025)
- Mohri, H. and Yasumasu, I.** (1963). Studies on the respiration of sea-urchin spermatozoa. V. The effects of P_{CO2}. *J. Exp. Biol.* **40**, 573-586.
- Morisawa, M.** (1985). Initiation mechanism of sperm motility at spawnig in teleosts. *Zool. Sci.* **2**, 605-615.
- Morisawa, M. and Suzuki, K.** (1980). Osmolality and potassium ion: their roles in initiation of sperm motility in teleosts. *Sci.* **210**, 1145-1147. (doi: 10.1126/science.7444445)
- Nakajima, A., Morita, M., Takemura, A., Kamimura, S. and Okuno, M.** (2005). Increase in intracellular pH induces phosphorylation of axonemal proteins for activation of flagellar motility in starfish sperm. *J. Exp. Biol.* **208**, 4411-4418. (doi: 10.1242/jeb.01906)
- Ohta, H., Kawamoto, T., Isowa, K., Aoki, H., Hayashi, M., Narita, T. and Komaru, A.** (2007). Motility of spermatozoa obtained from testes of Japanese pearl oyster *Pinctada fucata martensii*. *Fish. Sci.* **73**, 107-111. (doi: 10.1111/j.1444-2906.2007.01308.x)
- Paniagua-Chavez, C.G., Buchanan, J.T. and Tiersch, T.R.** (1998). Effect of extender solutions and dilution on motility and fertilizing ability of Eastern oyster sperm. *J. Shellfish Res.* **17**, 231-237.
- Phelps, H. and Warner, K.A.** (1990). Estuarine sediment bio-assay with oyster pediveliger larvae (*Crassostrea gigas*). *Bull. Environ. Contam. Toxicol.* **44**, 197-204. (doi: 10.1007/BF01700136)

- Piferrer, F., Beaumont, A., Falguiere, J.C., Flajshans, M., Haffray, P. and Colombo, L.** (2009). Polyploid fish and shellfish: production, biology and applications to aquaculture for performance improvement and genetic containment. *Aquaculture* **293**, 125–156. (doi.org/10.1016/j.aquaculture.2009.04.036).
- Robert, R. and Gérard, A.** (1999). Bivalve hatchery technology: the current situation for the Pacific oyster *Crassostrea gigas* and the scallop *Pecten maximus* in France. *Aquat. Living Resour.* **12**, 121–130. (doi: 10.1016/S0990-7440(99)80021-7).
- Ross, P.M., Parker, L., O’Conner, W.A. and Bailey, E.A.** (2011). The impact of Ocean Acidification on reproduction, Early Development and Settlement of Marine Organisms. *Water* **3**, 1005-1030. (doi: 10.3390/w3041005)
- Rothschild, L.** (1928). The physiology of sea urchin spermatozoa. Senescence and the dilution effect. *J. Exp. Biol.* **25**, 353-368.
- Schackmann, R.W., Christen, R. and Shapiro, B.M.** (1981). Membrane potential depolarisation and increased intracellular pH accompany the acrosome reaction of sea urchin sperm. *Proc. Natl. Acad. Sci. USA* **78**, 6066-6070.
- Schlegel, P., Havenhand, J.N., Gillings, M.R. and Williamson, J.E.** (2012). Individual variability in reproductive success determines winners and losers under ocean acidification: a case study with sea urchin. *Plos one* **7**, e53118. (doi: 10.1371/journal.pone.0053118)
- Steele, S. and Mulcahy, M.** (1999). Gametogenesis of the oyster *Crassostrea gigas* in southern Ireland. *J. Mar. Biol. Assoc. U.K.* **79**, 676-686. (doi: 10.1017/S0025315498000836)
- Suquet, M., Billard, R., Cosson, J., Dorange, G., Chauvaud, L., Mugnier, C. and Fauvel, C.** (1994). Sperm features in turbot (*Scophthalmus maximus*): a comparison with other freshwater and marine fish species. *Aquat. Living Resour.* **7**, 283-294. (doi: 10.1051/alr:1994031)
- Suquet, M., Malo, F., Quéau, I., Ratiskol, D., Quéré, C., Le Grand, J. and Fauvel, C.** (2016). Seasonal variation of sperm quality in Pacific oyster (*Crassostrea gigas*). *Aquaculture* **464**, 638-641. (doi :10.1016/j.aquaculture.2016.07.016)
- Vihtakari, M., Hendriks, I.E., Holding, J., Renaud, P.E., Duarte, C.M. and Havenhand, J.N.** (2013). Effects of ocean acidification and warming on sperm activity and early life stages of the Mediterranean mussel (*Mytilus galloprovincialis*). *Water* **5**, 1890-1915. (doi: 10.3390/w5041890)

- Wilson-Leedy, J.G. and Ingermann, R.L.** (2007) Development of a novel CASA system baed on open source software for characterization of zebrafish sperm motility parameters. *Theriogenol.* **67**, 661-672. (doi: 10.1016/j.theriogenology.2006.10.003)
- Zhao, X., Yu, H., Kong, L. and Li, Q.** (2012). Transcriptomic responses to salinity stress in the Pacific oyster *Crassostrea gigas*. *Plos one* **7**, e46244. (doi: 10.1371/journal.pone.0046244)

Figures




Fig. 1. Effects of salinity on percentage of motile spermatozoa and VAP after 1 and 5 min after activation. Motile spermatozoa (A, $P = 0.000$), VAP (B, $P = 0.003$), VAP: Velocity of the Average Path. VAP in salinities of 3.9 ‰, 8.9 ‰ (1 and 5 min after activation), and 14 ‰ (1 min after activation) were not measured because of low number of motile spermatozoa. VAP was not significantly different among salinities for spermatozoa activated for 1 min. Different letters refer to significantly different results (mean \pm s.e.m., lower case letters: 1 min after sperm activation, upper case letters: 5 min after sperm activation, $n = 5$ oysters). Data were compared using one way ANOVA and Fisher a posteriori test.




Fig. 2. Effects of pH on percentage of motile spermatozoa and VAP after 1 and 5 min after activation. Motile spermatozoa (A, $P = 0.000$), VAP (B, $P = 0.000$), VAP: Velocity of the Average Path. Different letters refer to significantly different results (mean \pm s.e.m., lower case letters: 1 min after sperm activation, upper case letters: 5 min after sperm activation, $n = 5$ oysters). Data were compared using one way ANOVA and Fisher a posteriori test.




Fig. 3. Effect of NH₄Cl (0 or 30 mM) on percentage of motile spermatozoa and VAP after 1 and 5 min after activation. Motile spermatozoa (A, $P = 0.000$), VAP (B), VAP: Velocity of the Average Path. VAP was not significantly different for both NH₄Cl concentrations (B). Different letters refer to significantly different results (mean \pm s.e.m., lower case letters: 1 min after sperm activation, upper case letters: 5 min after sperm activation, $n = 5$ oysters). Data were compared using one way ANOVA and Fisher a posteriori test.




Fig. 4. Effects of ion deprivation on percentage of motile spermatozoa and VAP after 1 and 5 min after activation. Motile spermatozoa (A, $P = 0.006$ at 1 min and $P = 0.000$ at 5 min after activation), VAP (B), VAP: Velocity of the Average Path. VAP in Na^+ -free ASW was not measured because of low number of motile spermatozoa. No movement of spermatozoa was observed in ion-free ASW. Sperm VAP was not significantly affected by a lack of Ca^{2+} , K^+ or Mg^{2+} in ASW. ASW: artificial seawater; different letters refer to significantly different results (mean \pm s.e.m., lower case letters: 1 min after sperm activation, upper case letters: 5 min after sperm activation, $n = 5$ oysters). Data were compared using one way ANOVA and Fisher a posteriori test.

Tables

Table 1. Chemical composition of activating media

Activating medium	Chemical composition of activating medium
ASW	516 mM NaCl, 10.4 mM KCl, 34 mM MgCl ₂ 2H ₂ O, 22 mM MgSO ₄ 7H ₂ O, 11 mM CaCl ₂ 2H ₂ O
Na ⁺ -free ASW	ASW without NaCl, NaCl of ASW substituted by adding 516mM choline chloride
Ca ²⁺ -free ASW	ASW without CaCl ₂ 2H ₂ O* 1mM EGTA, chelating Ca ²⁺
K ⁺ -free ASW	ASW without KCl *
Mg ²⁺ -free ASW	ASW without MgCl ₂ 2H ₂ O and MgSO ₄ 7H ₂ O MgCl ₂ 2H ₂ O substituted by adding 56mM choline chloride 1mM EDTA, chelating Mg ²⁺
Ion-free ASW	1M solution of glucose diluted in MilliQ water 1mM EGTA

ASW: artificial seawater, EGTA: ethylene glycol-bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid, EDTA: Ethylenediaminetetraacetic acid

* concentration of the ion studied in ASW is low and did not need to be substituted using choline chloride to maintain ASW osmotic pressure

Table 2. Composition of seminal fluid of the Pacific oyster

Parameter of the seminal fluid	Mean ± s.e.m.
Osmotic pressure (mosmol/L)	1061 ± 6
pH	5.82 ± 0.22
Cl ⁻ (mM)	521.3 ± 4.9
Na ⁺ (mM)	426.9 ± 4.1
K ⁺ (mM)	32.3 ± 1.4
Mg ²⁺ (mM)	50.4 ± 0.5
SO ₄ ²⁻ (mM)	25.5 ± 0.4
Ca ²⁺ (mM)	9.4 ± 0.1

n = 22 oysters, except n = 3 for osmotic pressure

Table 3. Composition of seminal fluid in some invertebrate species

Species	pH	Na ⁺ (mM)	K ⁺ (mM)	Mg ²⁺ (mM)	Ca ²⁺ (mM)	Reference
Pacific oyster <i>Crassostrea gigas</i>	5.5 - 6.5	360 ± 14	18 ± 1		11 ± 0	Alavi et al., 2014
Japanese scallop <i>Pactinopecten yessoensis</i>	7.0 ± 0.2	351 ± 83	41 ± 1		7 ± 1	Alavi et al., 2014
Sea urchin <i>Lytechinus pictus</i>	7.5 - 7.6	440	19	48	10	Johnson et al., 1983
Sea urchin <i>Strongylocentrotus purpuratus</i>	7.7	428	27		10	Christen et al., 1986

mean ± s.e.m.




Fig. S1. Percentage of motile spermatozoa and VAP in seawater and artificial seawater after 1 and 5 min after activation. Motile (A), VAP (B), VAP: Velocity of the Average Path, SW: seawater, ASW: artificial seawater, mean \pm SEM, n = 5 oysters.




Fig. S1. Percentage of motile spermatozoa and VAP in seawater and artificial seawater after 1 and 5 min after activation. Motile (A), VAP (B), VAP: Velocity of the Average Path, SW: seawater, ASW: artificial seawater, mean \pm SEM, n = 5 oysters.