

In vivo genetic manipulation of cortical progenitors in gyrencephalic carnivores using *in utero* electroporation

Hiroshi Kawasaki^{1,2,*}, Tomohisa Toda^{1,2} and Kaori Tanno^{1,2}

¹Department of Molecular and Systems Neurobiology, Graduate School of Medicine, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

²Global COE Program "Comprehensive Center of Education and Research for Chemical Biology of the Diseases", The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

*Author for correspondence (hiroshi-kawasaki@umin.ac.jp)

Biology Open 2, 95–100

doi: 10.1242/bio.20123160

Received 22nd September 2012

Accepted 7th November 2012

Summary

Brain structures such as the outer subventricular zone (OSVZ) and the inner fiber layer (IFL) in the developing cerebral cortex are especially prominent in higher mammals. However, the molecular mechanisms underlying the formation of the OSVZ are still largely unknown, mainly because genetic manipulations that can be applied to the OSVZ in higher mammals had been poorly available. Here we developed and validated a rapid and efficient genetic manipulation technique for germinal zones including the OSVZ using *in utero* electroporation in developing gyrencephalic carnivore ferrets. We also determined the optimal conditions for using *in utero* electroporation to express transgenes in germinal zones. Using our electroporation procedure, the morphology of GFP-positive cells in the OSVZ was clearly visible even without immunostaining, and multiple genes were efficiently

co-expressed in the same cells. Furthermore, we uncovered that fibers, which seemed to correspond to those in the IFL of monkeys, also existed in ferrets, and were derived from newly generated cortical neurons. Our technique promises to be a powerful tool for investigating the fundamental mechanisms underlying the formation and abnormalities of the cerebral cortex in higher mammals.

© 2012. Published by The Company of Biologists Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial Share Alike License (<http://creativecommons.org/licenses/by-nc-sa/3.0>).

Key words: Ferret, Cerebral cortex, *In utero* electroporation, Outer subventricular zone, Inner fiber layer, Outer radial glia

Introduction

Expansion and folding of the cerebral cortex are the most notable features of higher mammals such as carnivores and primates. A major underlying cause of these features seems to be the increase in population size of neural progenitors in the outer subventricular zone (OSVZ), a specialized germinal zone characteristic of the developing cerebral cortex in higher mammals. To better understand the OSVZ, it is therefore important to uncover the mechanisms regulating the differentiation and proliferation of neural progenitors in the cerebral cortex of developing higher mammals (Dehay and Kennedy, 2007; Rakic, 2009; Fietz and Huttner, 2011; Lui et al., 2011; Borrell and Reillo, 2012; Hevner and Haydar, 2012; Molnár and Clowry, 2012).

Cortical neurons arise from radial glial cells (RG cells, also known as apical progenitors/apical RG/ventricular RG), the epithelial stem cells that line the cerebral ventricles and extend apical fibers and basal fibers (Malatesta et al., 2000; Miyata et al., 2001; Noctor et al., 2001). RG cells in the ventricular zone (VZ) undergo multiple rounds of asymmetric divisions and generate intermediate progenitor cells (IP cells/basal progenitors) that migrate into the subventricular zone (SVZ) and further proliferate to increase neuronal number (Haubensak et al., 2004; Noctor et al., 2004). Corticogenesis in higher mammals is distinguished by the appearance of the large SVZ that has an inner (ISVZ) and

outer region (OSVZ), often split by a thin layer of fibers, the inner fiber layer (IFL) (Smart et al., 2002; Zecevic et al., 2005). Recent studies identified a novel class of progenitor cells found in the OSVZ, termed OSVZ radial glial cells (oRG cells, also known as outer RG/basal RG/intermediate RG/translocating RG). Unlike RG cells in the VZ, oRG cells are unipolar, with a basal fiber that ascends toward the pia without an apical fiber that descends toward the ventricle (Fietz et al., 2010; Hansen et al., 2010).

Although there have been extensive anatomical and histochemical investigations of the OSVZ, and though previous pioneering studies uncovered the involvement of the Notch signaling and β 3-integrin in the development of the OSVZ using cultured slices *in vitro* (Fietz et al., 2010; Hansen et al., 2010), a molecular understanding of the OSVZ *in vivo* is still limited. This is mainly because rapid and efficient genetic manipulations that can be applied to germinal zones of the cerebral cortex in higher mammals had been poorly available. *In utero* electroporation is a useful technique to express genes of interest in the living rodent brain (Fukuchi-Shimogori and Grove, 2001; Saito and Nakatsuji, 2001; Tabata and Nakajima, 2001; Sehara et al., 2010; Ako et al., 2011), but successful application of *in utero* electroporation in higher mammals had not been achieved. Therefore, we recently developed and validated a rapid and efficient procedure for *in*

utero electroporation in gyrencephalic carnivore ferrets and demonstrated that genes of interest can be efficiently expressed in post-mitotic neurons (Kawasaki et al., 2012). Here we show that transgenes can be also efficiently expressed in neural progenitors in germinal zones including the OSVZ using *in utero* electroporation in the developing ferret. We also determined the optimal conditions for using *in utero* electroporation to express transgenes in germinal zones. Furthermore, using our electroporation procedure, we uncovered that fibers, which seemed to correspond to those in the IFL of monkeys, also existed in ferrets, and were derived from newly generated cortical neurons. Our method promises to be a powerful tool for investigating the molecular mechanisms underlying the development of the OSVZ in higher mammals. Uncovering the developmental processes of the OSVZ using higher mammals would help lead to the ultimate goal of understanding the human brain and its diseases.

Materials and Methods

Animals

Normally pigmented, sable ferrets (*Mustela putorius furo*) were purchased from Marshall Farms (North Rose, NY). Ferrets were maintained as described previously (Kawasaki et al., 2004; Iwai and Kawasaki, 2009; Iwai et al., 2012; Kawasaki et al., 2012). The day of birth was counted as postnatal day 0 (P0). All procedures were performed in accordance with a protocol approved by the University of Tokyo Animal Care Committee.

In utero electroporation procedure for ferrets

By modifying the procedure for *in utero* electroporation of rodents, we recently created a procedure for *in utero* electroporation to express transgenes in post-mitotic neurons of the ferret cerebral cortex (Kawasaki et al., 2012). Briefly, pregnant ferrets were anesthetized with sodium pentobarbital, and their body temperature was monitored and maintained using a heating pad. The uterine horns were exposed and kept wet by adding drops of PBS intermittently. The location of embryos was visualized with transmitted light delivered through an optical fiber cable. The pigmented iris was visible, and this enabled us to assume the location of the lateral ventricle. Approximately 2–5 μ l of DNA solution was injected into the lateral ventricle at the indicated ages using a pulled glass micropipette. Because the position and shape of the placenta in ferrets are more obscure compared with those in mice, care should be taken not to damage the placenta with glass micropipettes. Each embryo within the uterus was placed between tweezer-type electrodes with a diameter of 5 mm (CUY650-P5; NEPA Gene, Japan). Square electric pulses (50–150 V, 50 ms) were passed 5 times at 1-second intervals using an electroporator (ECM830, BTX). Care was taken to quickly place embryos back into the abdominal cavity to avoid excessive temperature loss. The wall and skin of the abdominal cavity were sutured, and the embryos were allowed to develop normally.

Plasmids

pCAG-GFP and pCAG-mCherry were described previously (Sehara et al., 2010). Plasmids were purified using the Endofree plasmid Maxi kit (Qiagen, Valencia, CA). Prior to *in utero* electroporation procedures, plasmid DNA was diluted to 0.5–3.0 mg/ml in 1 \times PBS, and Fast Green solution was added to a final concentration of 0.5% to monitor the injection. For co-transfection, a mixture of pCAG-GFP and pCAG-mCherry was used.

Preparation of sections

Preparation of sections was performed as described previously with slight modifications (Matsubayashi et al., 2008; Toda et al., 2008). Briefly, ferrets were deeply anesthetized with pentobarbital and transcardially perfused with 4% paraformaldehyde (PFA), and then the brain was dissected. Alternatively, brains were taken from deeply anesthetized ferrets, and immersion-fixation was performed using 4% PFA. Then, the brains were cryoprotected by overnight immersion in 30% sucrose and embedded in OCT compound. Sections of 14–50 μ m thickness were incubated with 1 μ g/ml Hoechst 33342, washed and mounted. Experiments were repeated at least three times and gave consistent results.

Immunohistochemistry

Immunohistochemistry was performed as described previously with slight modifications (Matsubayashi et al., 2008; Hayakawa and Kawasaki, 2010).

Sections were made using a cryostat, permeabilized with 0.1–0.5% Triton X-100/PBS, and incubated overnight with primary antibodies, which included anti-Sox2 antibody (R&D Systems), anti-Pax6 antibody (Covance), anti-Tbr2 antibody (Abcam) and anti-neurofilament-M antibody (Chemicon). After incubation with secondary antibodies and Hoechst 33342, the sections were washed and mounted. Experiments were repeated at least three times in different animals and gave consistent results.

Results

Efficient expression of GFP in germinal zones of the ferret cerebral cortex using *in utero* electroporation

We recently established and reported *in utero* electroporation procedure for ferrets and successfully expressed GFP in post-mitotic neurons of the cerebral cortex (Kawasaki et al., 2012). We therefore assumed that our procedure is also applicable to manipulation of gene expression in the OSVZ. To identify the appropriate time point for performing *in utero* electroporation to express transgenes in germinal zones, we carried out *in utero* electroporation at various time points during development. Because the OSVZ is prominent soon after birth in ferrets (Fietz et al., 2010; Reillo and Borrell, 2012), we dissected ferret babies at postnatal day 0 (P0). Consistent with our previous report (Kawasaki et al., 2012), GFP fluorescence was clearly visible on the brain surface (Fig. 1A, arrowhead), and numerous GFP-




Fig. 1. GFP expression in ferrets induced by *in utero* electroporation performed at various time points during development. (A) *In utero* electroporation was performed at E35, and GFP fluorescence was examined at P0. A dorsal view of the ferret brain is shown. GFP fluorescence was clearly visible in the cerebral cortex (arrowhead). a, anterior; p, posterior. (B) GFP-positive cells in the cerebral cortex. A coronal section is shown. GFP-positive cells were clearly visible even without immunostaining. *Lateral ventricle. (C) *In utero* electroporation was carried out at the indicated time points during development. Coronal sections were prepared at P0 and were stained with Hoechst 33342. Magnified images of the cerebral cortex are shown. When electroporation was performed at E31, most GFP-positive cells had already moved into the cortical plate at P0 (arrowhead). When electroporation was carried out at E40, many GFP-positive cells were found in germinal zones (arrow). CP, cortical plate; OSVZ, outer subventricular zone; ISVZ, inner subventricular zone; VZ, ventricular zone. Scale bars: 2 mm (A), 1 mm (B) and 200 μ m (C).

positive cells were observed in the cerebral cortex (Fig. 1B), indicating that cortical cells were efficiently transfected with GFP using our *in utero* electroporation procedure in ferrets.

When ferrets were electroporated with pCAG-GFP at E31, we found that most GFP-positive cells had already moved into the cortical plate at birth (Fig. 1C, E31, arrowhead). Electroporation at E35 resulted in GFP-positive cells distributed throughout the developing cortex including the OSVZ (Fig. 1C, E35). When electroporation was performed at E40, germinal zones were prominently labeled with GFP (Fig. 1C, E40, arrow). These results suggest that our electroporation procedure is applicable for expressing transgenes in neural progenitors and that *in utero* electroporation between E35 and E40 is appropriate for obtaining transfected neural progenitors at birth.

Germinal zones in the cerebral cortex of developing ferrets can be distinguished by the expression patterns of progenitor marker molecules such as Sox2, Pax6 and Tbr2. We therefore compared the distribution pattern of GFP-positive cells with those of progenitor markers using immunohistochemistry. We performed *in utero* electroporation at E37, and sections prepared at E40 were stained with anti-Pax6, Sox2 and Tbr2 antibodies (Fig. 2). Pax6 and Sox2 are strongly expressed in the VZ, whereas Tbr2 is preferentially expressed in the ISVZ (Fietz et al., 2010; Reillo and Borrell, 2012; Martínez-Cerdeño et al., 2012). Our immunohistochemical studies showed that many GFP-positive cells were located not only in the ISVZ but also in the VZ and the OSVZ (Fig. 2). These results clearly indicate that our electroporation procedure is applicable to manipulation of gene expression in germinal zones of the developing ferret cerebral cortex.

Visualization of the morphology of GFP-positive cells in the OSVZ

Because the morphology of neural progenitors changes during development, it would be useful if our electroporation procedure




Fig. 2. Distribution of GFP-positive cells and progenitor markers. *In utero* electroporation was performed at E37, and sections were prepared at E40. The sections were immunostained with anti-Sox2, anti-Pax6 and anti-Tbr2 antibodies. The cerebral cortex is shown in A. The areas within the white boxes are magnified and are shown in B. CP, cortical plate; OSVZ, outer subventricular zone; ISVZ, inner subventricular zone; VZ, ventricular zone. Scale bars: 200 μ m (A) and 100 μ m (B).

were applicable for visualizing the morphology of single cells in the cerebral cortex of developing ferrets. We therefore examined whether GFP expression was strong enough for examining the morphology of cells without immunostaining. High magnification images clearly demonstrated that the morphology of individual GFP-positive cells in the OSVZ was clearly visible (Fig. 3C). When sections were immunostained with anti-Sox2 antibody, we found GFP-positive cells that were also positive for Sox2 in the OSVZ (Fig. 3D, arrowhead), suggesting that these GFP-positive cells are oRG cells. Consistently, these GFP/Sox2 double-positive cells extended their processes toward the pial surface without having apical fibers (Fig. 3D, arrowhead). Anti-Pax6 antibody gave results similar to those obtained with anti-Sox2 antibody (Fig. 3E, arrowhead). Combining our electroporation procedure and time-lapse imaging would contribute to uncovering the dynamics of the morphological changes and developmental lineages of progenitor cells in higher mammals during development.




Fig. 3. The morphology of GFP-positive cells in the OSVZ. *In utero* electroporation was performed at E35, and coronal sections were prepared at P0 and were stained with Hoechst 33342. Many GFP-positive cells were distributed throughout the cortex (A). The areas within the white boxes in A are magnified and are shown in B. The areas within the white boxes in B are shown in C. Note that the morphology of GFP-positive cells was clearly visible even without immunostaining. (D,E) The sections were immunostained with anti-Sox2 antibody (D) and anti-Pax6 antibody (E), and high magnification images of the OSVZ are shown. The GFP-positive cells (arrowheads) expressed Sox2 (D) and Pax6 (E), and had basal fibers but not apical fibers, suggesting that these cells are oRG cells. GFP-positive fibers running tangentially in the inner OSVZ were also visible (arrow). *Lateral ventricle. CP, cortical plate; OSVZ, outer subventricular zone; ISVZ, inner subventricular zone; VZ, ventricular zone. Scale bars: 1 mm (A), 500 μ m (B), 100 μ m (C) and 20 μ m (D,E).

Co-expression of GFP and mCherry in germinal zones of the ferret cerebral cortex

To investigate the molecular mechanisms underlying the formation of the OSVZ, it would be useful if multiple genes, such as GFP plus genes of interest, could be co-transfected in progenitor cells of ferrets. We therefore examined co-transfection efficiency by transfecting a mixture of pCAG-GFP and pCAG-mCherry at E35 and dissected the brain at birth. As we have shown, numerous GFP-positive cells were distributed throughout the cortex (Fig. 4A). We then examined co-localization of GFP and mCherry at the cellular level using high magnification images. We found that most of GFP-positive cells were also positive for mCherry ($91.8 \pm 5.7\%$, 6 sections from 3 animals) (Fig. 4B). These results indicate that the co-transfection efficiency in the OSVZ is reasonably high.

Distribution of GFP-positive cells during development

Using *in utero* electroporation, we next examined changes in the distribution of GFP-positive cells during development in ferrets. We carried out *in utero* electroporation at E35–37 and dissected ferret babies at E40, P2 and P10. When ferret babies were dissected at E40, GFP-positive cells were distributed in the VZ, ISVZ and OSVZ (Fig. 5A,B). Basal fibers that ascend toward the pia were clearly visible (Fig. 5A,B, arrowheads). GFP-positive cells were located throughout the cortex at P2 and found mostly in the cortical plate at P10 (Fig. 5A), suggesting that most GFP-positive cells are migrating at P2, and the migration has been mostly completed by P10.

Identification and characterization of IFL-like fibers in ferrets

Although the IFL, which separates the ISVZ and the OSVZ, is one of the prominent features of the cerebral cortex of monkeys and humans (Smart et al., 2002; Zecevic et al., 2005), a previous study reported that an obvious fiber layer between the ISVZ and the OSVZ was difficult to distinguish in ferrets (Martínez-Cerdeño et al., 2012). It seemed possible that the IFL is specific to primates, and ferrets do not have fibers corresponding to those found in the IFL of monkeys. Conversely, it also seemed possible that ferrets do have fibers similar to those in the IFL of monkeys, but the fibers do not form a specific cytoarchitectonic layer and run within the neighboring OSVZ and/or ISVZ. Interestingly, we found GFP-positive fibers running tangentially in the OSVZ (Fig. 5A, P2 and P10, arrows; Fig. 5B, P2, square bracket. See also Fig. 3D, arrow). The GFP-positive fibers were predominantly located in the lower OSVZ (Fig. 5, P2 and P10).




Fig. 4. Double labeling of OSVZ cells using *in utero* electroporation. Cortical cells were co-transfected with pCAG-GFP and pCAG mCherry using *in utero* electroporation at E35. The brain was dissected out at P0, and coronal sections were made. (A) Numerous GFP-positive cells and mCherry-positive cells were distributed throughout the cortex. The areas within the white boxes, which correspond to the OSVZ, were magnified and are shown in B. Note that most GFP-positive cells in the OSVZ were also positive for mCherry. Scale bars: 1 mm (A) and 200 μm (B).




Fig. 5. Distribution patterns of GFP-positive cells and IFL-like fibers during development. *In utero* electroporation was performed at E35–37, and the distribution patterns of GFP-positive cells and IFL-like fibers in the cerebral cortex were examined at the indicated time points during development. The coronal sections of the cerebral cortex were stained with Hoechst 33342. The areas in the white boxes were magnified and are shown in B. At E40, GFP-positive cells were mainly found in germinal zones. Note that basal fibers that ascend toward the pia are clearly visible (arrowheads in A and B). GFP-positive cells were distributed throughout the cortex at P2 and were mostly found in the cortical plate at P10. GFP-positive IFL-like fibers were found preferentially in the lower OSVZ (arrows in A and square bracket in B). CP, cortical plate; OSVZ, outer subventricular zone; ISVZ, inner subventricular zone; VZ, ventricular zone. Scale bars: 500 μm (A, P2 and P10), 200 μm (A, E40; B, P2) and 100 μm (B, E40).

This result is consistent with the idea that fibers, which correspond to those in the IFL of monkeys, also exist in ferrets and are running mainly in the lower OSVZ, although it is important to investigate whether the GFP-positive fibers found in this study are indeed homologous to fibers in the IFL of monkeys and humans. A previous study pointed out that the ferret OSVZ could be further divided into inner and outer subdivisions by Nissl staining, and the inner subdivision of the OSVZ had some cytoarchitectonic features similar to the IFL in monkeys (Reillo and Borrell, 2012). This inner subdivision of the OSVZ seems to correspond to the lower OSVZ, where we found the GFP-positive fibers.

Although the IFL is a prominent feature of the cerebral cortex in higher mammals (Smart et al., 2002; Zecevic et al., 2005), the origin of the IFL was unclear (Molnár and Clowry, 2012). One possibility was that the IFL consists of the fibers derived from newly generated cortical neurons, while it was also possible that early thalamocortical projections constitute the IFL (Molnár and Clowry, 2012). Because we expressed GFP in cortical cells but not in the thalamus, our results indicate that IFL-like fibers in ferrets consists of, at least partially, fibers derived from newly generated cortical neurons and/or neural progenitors. Interestingly, at E40, when GFP expression was restricted to neural progenitors, IFL-like fibers were invisible (Fig. 5, E40). This result suggests cortical neurons rather than neural progenitors are responsible for IFL-like fibers in the developing ferret cortex. Consistently, neurofilament-M was found

in IFL-like fibers (Fig. 6). Because both the IFL and the increased number of cortical neurons are features of higher mammals, it seems plausible that the fibers in the IFL are derived from evolutionarily new cortical neurons that were added through cortical expansion during evolution. In addition, because the IFL is prominent in higher mammals, it would be intriguing to uncover its functional role.

Discussion

Here we have shown that transgenes can be efficiently expressed in neural progenitors in developing ferrets. We also determined the optimal experimental conditions to express transgenes in neural progenitors. Our procedure should provide a rapid means to investigate the molecular mechanisms underlying the development of the cerebral cortex in higher mammals. In addition, our results suggest that IFL-like fibers also exist in ferrets and are derived from newly generated cortical neurons.

Development of the cerebral cortex in higher mammals

Corticogenesis in higher mammals is characterized by the appearance of the OSVZ and the IFL. Earlier studies suggested that the proliferation of cells within the OSVZ contribute to the expansion of the cerebral cortex in higher mammals. Although anatomical and histochemical investigations have been performed extensively, the molecular mechanisms underlying the formation of the OSVZ are still unclear. Because previous pioneering studies reported the involvement of the Notch signaling and β 3-integrin in the development of oRG cells using cultured slices *in vitro* (Fietz et al., 2010; Hansen et al., 2010), it would be important to examine the roles of the Notch signaling and β 3-integrin *in vivo* using our method. Our method promises to be a powerful tool for investigating the molecular mechanisms underlying the proliferation and differentiation of neural progenitors in higher mammals.

Importantly, our results showed that the expression level of GFP was high enough to examine the morphology of transfected cells without using immunostaining. Therefore, it seems plausible that morphological changes of progenitors can be examined in living ferret neonates. Because RG cells, IP cells and oRG cells have distinct characteristic morphologies, it would be intriguing to examine the morphological transitions of these cells *in vivo*. It would also be interesting to investigate other features *in vivo* such as the cleavage plane during mitosis and cellular locomotion such as interkinetic nuclear migration and mitotic somal translocation.




Fig. 6. Expression of neurofilament-M in IFL-like fibers. Sections were prepared at E40 and stained with anti-neurofilament-M antibody. Confocal images showed that GFP-positive IFL-like fibers were also positive for neurofilament-M (NF-M). Magnified images of the areas within the white boxes are shown on the right (A,B). Scale bar: 20 μ m.

The advantages of ferrets

Usually more than 6 ferret babies are born from one pregnant mother. This large number of babies per pregnant mother relative to other higher mammals such as the marmoset is an important advantage of ferrets. This enables us to examine various experimental conditions and to obtain a sufficient number of experimental samples.

Brain structures such as the OSVZ, the IFL, ocular dominance columns (ODCs) and the parallel visual pathways (i.e. the magnocellular, parvocellular and koniocellular pathways) are especially prominent in higher mammals including ferrets. Although these structures have been believed to be important, the molecular mechanisms underlying the formation of these structures and their functional significance are still elusive. This is mainly because rapid and efficient genetic methods for manipulating genes in the brains of higher mammals had been poorly available. Therefore, our *in utero* electroporation procedure for ferrets should open the door to the next generation of neuroscientific experiments using higher mammals.

Recently, several groups including us have reported molecules expressed in the structures mentioned above that are unique to higher mammals (Kawasaki et al., 2004; Murray et al., 2008; Johnson et al., 2009; Yamamori, 2011; Bernard et al., 2012; Iwai et al., 2012). For example, we uncovered that the Forkhead transcription factor FoxP2 was selectively expressed in the parvocellular pathway in the dLGN of ferrets and monkeys (Iwai et al., 2012). Furthermore, recent studies provided lists of genes expressed in the VZ and the SVZ of mice and various cortical regions in monkeys (Ayoub et al., 2011; Bernard et al., 2012). Combining these molecules and our electroporation procedure, gain-of-function and loss-of-function studies using higher mammals are now ready to be conducted.

Advantages of *in utero* electroporation

It seems plausible to make transgenic ferrets using virus vectors because the successful application of virus vectors to make transgenic monkeys and marmosets was reported (Chan et al., 2001; Lois et al., 2002; Sasaki et al., 2009). However, compared with virus vectors, *in utero* electroporation has several advantageous features. First, it does not take a long time to obtain transfected animals. Transfected ferrets should be available within a couple of days. Second, multiple genes can easily be introduced simultaneously, as shown in this study. Co-transfection of GFP plus genes of interest should be useful. Third, transgenes can be selectively expressed in appropriate brain regions, even without using specific promoters, by modifying the direction of electrodes and the age when electroporation is performed. Finally, if necessary, cell type-specific promoters can be utilized. Because germinal zones of the cerebral cortex contain a heterogeneous population of progenitor cells (Hansen et al., 2010; Reillo and Borrell, 2012), it might be important to express transgenes in a cell type-specific manner to investigate the mechanisms underlying development of each progenitor. We recently reported that *in utero* electroporation is useful for exploring promoter regions and found that the newly generated Thy1S promoter resulted in sparse neuronal labeling in the cerebral cortex and the hippocampus of mice (Ako et al., 2011). Thus, our *in utero* electroporation procedure for ferrets described here could contribute to identifying cell type-specific promoters for each progenitor.

Theoretically, using our *in utero* electroporation procedure, it should be possible to express transgenes in neural progenitors

including shRNA constructs, optogenetic molecules (e.g. channelrhodopsin and halorhodopsin), neuronal activity reporters (e.g. GFP-based Ca²⁺ sensors) and activity-modifying channels (e.g. Kir2.1 and NaChBac) (Boyden et al., 2005; Deisseroth et al., 2006; Arenkiel et al., 2007; Gradinaru et al., 2010). Combining these molecules with *in utero* electroporation in ferrets would contribute toward an understanding of the formation, evolution and function of the gyrencephalic brains of higher mammals. In addition, because our results indicate that *in utero* electroporation can be used not only in rodents but also in ferrets, it seems reasonable to speculate that *in utero* electroporation is applicable to other higher mammals such as primates. It would be intriguing to establish *in utero* electroporation protocols for primates.

Acknowledgements

We are grateful for Drs Shoji Tsuji, Haruhiko Bito, Takashi Kadowaki (The University of Tokyo), Eisuke Nishida (Kyoto University), Yoshiki Sasai (RIKEN-CDB) and Shigetada Nakanishi (Osaka Bioscience Institute) for their continuous encouragement. We thank Zachary Blalock and Kawasaki lab members for their helpful discussion and support. This work was supported by the 21st Century COE Program “Center for Integrated Brain Medical Sciences” from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), the Global COE Program “Comprehensive Center of Education and Research for Chemical Biology of the Diseases” from MEXT, Grant-in-Aid for Scientific Research from MEXT, PRESTO from Japan Science and Technology Agency, and Human Frontier Science Program. This work was also supported by Takeda Science Foundation, Takeda Medical Research Foundation, Astellas Foundation for Research on Metabolic Disorders, the Life Science Foundation of Japan, the Kurata Memorial Hitachi Science and Technology Foundation, Mitsubishi Foundation, Fukuda Foundation for Medical Technology, Yamada Science Foundation, Hokuto Foundation, Daiichi-Sankyo Foundation of Life Sciences, Research Foundation for Opto-Science and Technology, and Santen Pharmaceutical.

Competing Interests

The authors have no competing interests to declare.

References

- Ako, R., Wakimoto, M., Ebisu, H., Tanno, K., Hira, R., Kasai, H., Matsuzaki, M. and Kawasaki, H. (2011). Simultaneous visualization of multiple neuronal properties with single-cell resolution in the living rodent brain. *Mol. Cell. Neurosci.* **48**, 246-257.
- Arenkiel, B. R., Peca, J., Davison, I. G., Feliciano, C., Deisseroth, K., Augustine, G. J., Ehlers, M. D. and Feng, G. (2007). *In vivo* light-induced activation of neural circuitry in transgenic mice expressing channelrhodopsin-2. *Neuron* **54**, 205-218.
- Ayoub, A. E., Oh, S., Xie, Y., Leng, J., Cotney, J., Dominguez, M. H., Noonan, J. P. and Rakic, P. (2011). Transcriptional programs in transient embryonic zones of the cerebral cortex defined by high-resolution mRNA sequencing. *Proc. Natl. Acad. Sci. USA* **108**, 14950-14955.
- Bernard, A., Lubbers, L. S., Tanis, K. Q., Luo, R., Podtelezchnikov, A. A., Finney, E. M., McWhorter, M. M. E., Serikawa, K., Lemon, T., Morgan, R. et al. (2012). Transcriptional architecture of the primate neocortex. *Neuron* **73**, 1083-1099.
- Borrell, V. and Reillo, I. (2012). Emerging roles of neural stem cells in cerebral cortex development and evolution. *Dev. Neurobiol.* **72**, 955-971.
- Boyden, E. S., Zhang, F., Bamberg, E., Nagel, G. and Deisseroth, K. (2005). Millisecond-timescale, genetically targeted optical control of neural activity. *Nat. Neurosci.* **8**, 1263-1268.
- Chan, A. W., Chong, K. Y., Martinovich, C., Simerly, C. and Schatten, G. (2001). Transgenic monkeys produced by retroviral gene transfer into mature oocytes. *Science* **291**, 309-312.
- Dehay, C. and Kennedy, H. (2007). Cell-cycle control and cortical development. *Nat. Rev. Neurosci.* **8**, 438-450.
- Deisseroth, K., Feng, G., Majewska, A. K., Miesenböck, G., Ting, A. and Schnitzer, M. J. (2006). Next-generation optical technologies for illuminating genetically targeted brain circuits. *J. Neurosci.* **26**, 10380-10386.
- Fietz, S. A. and Huttner, W. B. (2011). Cortical progenitor expansion, self-renewal and neurogenesis—a polarized perspective. *Curr. Opin. Neurobiol.* **21**, 23-35.
- Fietz, S. A., Kelava, I., Vogt, J., Wilsch-Bräuninger, M., Stenzel, D., Fish, J. L., Corbeil, D., Riehn, A., Distler, W., Nitsch, R. et al. (2010). OSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling. *Nat. Neurosci.* **13**, 690-699.
- Fukuchi-Shimogori, T. and Grove, E. A. (2001). Neocortex patterning by the secreted signaling molecule FGF8. *Science* **294**, 1071-1074.
- Gradinaru, V., Zhang, F., Ramakrishnan, C., Mattis, J., Prakash, R., Diester, I., Goshen, I., Thompson, K. R. and Deisseroth, K. (2010). Molecular and cellular approaches for diversifying and extending optogenetics. *Cell* **141**, 154-165.
- Hansen, D. V., Lui, J. H., Parker, P. R. and Kriegstein, A. R. (2010). Neurogenic radial glia in the outer subventricular zone of human neocortex. *Nature* **464**, 554-561.
- Haubensak, W., Attardo, A., Denk, W. and Huttner, W. B. (2004). Neurons arise in the basal neuroepithelium of the early mammalian telencephalon: a major site of neurogenesis. *Proc. Natl. Acad. Sci. USA* **101**, 3196-3201.
- Hayakawa, I. and Kawasaki, H. (2010). Rearrangement of retinogeniculate projection patterns after eye-specific segregation in mice. *PLoS ONE* **5**, e11001.
- Hevner, R. F. and Haydar, T. F. (2012). The (not necessarily) convoluted role of basal radial glia in cortical neurogenesis. *Cereb. Cortex* **22**, 465-468.
- Iwai, L. and Kawasaki, H. (2009). Molecular development of the lateral geniculate nucleus in the absence of retinal waves during the time of retinal axon eye-specific segregation. *Neuroscience* **159**, 1326-1337.
- Iwai, L., Ohashi, Y., van der List, D., Usrey, W. M., Miyashita, Y. and Kawasaki, H. (2012). FoxP2 is a parvocellular-specific transcription factor in the visual thalamus of monkeys and ferrets. *Cereb. Cortex*. [Epub ahead of print].
- Johnson, M. B., Kawasawa, Y. I., Mason, C. E., Krsnik, Z., Coppola, G., Bogdanović, D., Geschwind, D. H., Mane, S. M., State, M. W. and Sestan, N. (2009). Functional and evolutionary insights into human brain development through global transcriptome analysis. *Neuron* **62**, 494-509.
- Kawasaki, H., Crowley, J. C., Livesey, F. J. and Katz, L. C. (2004). Molecular organization of the ferret visual thalamus. *J. Neurosci.* **24**, 9962-9970.
- Kawasaki, H., Iwai, L. and Tanno, K. (2012). Rapid and efficient genetic manipulation of gyrencephalic carnivores using *in utero* electroporation. *Mol. Brain* **5**, 24.
- Lois, C., Hong, E. J., Pease, S., Brown, E. J. and Baltimore, D. (2002). Germline transmission and tissue-specific expression of transgenes delivered by lentiviral vectors. *Science* **295**, 868-872.
- Lui, J. H., Hansen, D. V. and Kriegstein, A. R. (2011). Development and evolution of the human neocortex. *Cell* **146**, 18-36.
- Malatesta, P., Hartfuss, E. and Götz, M. (2000). Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage. *Development* **127**, 5253-5263.
- Martínez-Cerdeño, V., Cunningham, C. L., Camacho, J., Antczak, J. L., Prakash, A. N., Cziep, M. E., Walker, A. I. and Noctor, S. C. (2012). Comparative analysis of the subventricular zone in rat, ferret and macaque: evidence for an outer subventricular zone in rodents. *PLoS ONE* **7**, e30178.
- Matsubayashi, Y., Iwai, L. and Kawasaki, H. (2008). Fluorescent double-labeling with carbocyanine neuronal tracing and immunohistochemistry using a cholesterol-specific detergent digitonin. *J. Neurosci. Methods* **174**, 71-81.
- Miyata, T., Kawaguchi, A., Okano, H. and Ogawa, M. (2001). Asymmetric inheritance of radial glial fibers by cortical neurons. *Neuron* **31**, 727-741.
- Molnár, Z. and Clowry, G. (2012). Cerebral cortical development in rodents and primates. *Prog. Brain Res.* **195**, 45-70.
- Murray, K. D., Rubin, C. M., Jones, E. G. and Chalupa, L. M. (2008). Molecular correlates of laminar differences in the macaque dorsal lateral geniculate nucleus. *J. Neurosci.* **28**, 12010-12022.
- Noctor, S. C., Flint, A. C., Weissman, T. A., Dammerman, R. S. and Kriegstein, A. R. (2001). Neurons derived from radial glial cells establish radial units in neocortex. *Nature* **409**, 714-720.
- Noctor, S. C., Martínez-Cerdeño, V., Ivic, L. and Kriegstein, A. R. (2004). Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. *Nat. Neurosci.* **7**, 136-144.
- Rakic, P. (2009). Evolution of the neocortex: a perspective from developmental biology. *Nat. Rev. Neurosci.* **10**, 724-735.
- Reillo, I. and Borrell, V. (2012). Germinal zones in the developing cerebral cortex of ferret: ontogeny, cell cycle kinetics, and diversity of progenitors. *Cereb. Cortex* **22**, 2039-2054.
- Saito, T. and Nakatsuji, N. (2001). Efficient gene transfer into the embryonic mouse brain using *in vivo* electroporation. *Dev. Biol.* **240**, 237-246.
- Sasaki, E., Suemizu, H., Shimada, A., Hanazawa, K., Oiwa, R., Kamioka, M., Tomioka, I., Sotomaru, Y., Hirakawa, R., Eto, T. et al. (2009). Generation of transgenic non-human primates with germline transmission. *Nature* **459**, 523-527.
- Sehara, K., Toda, T., Iwai, L., Wakimoto, M., Tanno, K., Matsubayashi, Y. and Kawasaki, H. (2010). Whisker-related axonal patterns and plasticity of layer 2/3 neurons in the mouse barrel cortex. *J. Neurosci.* **30**, 3082-3092.
- Smart, I. H., Dehay, C., Giroud, P., Berland, M. and Kennedy, H. (2002). Unique morphological features of the proliferative zones and postmitotic compartments of the neural epithelium giving rise to striate and extrastriate cortex in the monkey. *Cereb. Cortex* **12**, 37-53.
- Tabata, H. and Nakajima, K. (2001). Efficient *in utero* gene transfer system to the developing mouse brain using electroporation: visualization of neuronal migration in the developing cortex. *Neuroscience* **103**, 865-872.
- Toda, T., Hayakawa, I., Matsubayashi, Y., Tanaka, K., Ikenaka, K., Lu, Q. R. and Kawasaki, H. (2008). Termination of lesion-induced plasticity in the mouse barrel cortex in the absence of oligodendrocytes. *Mol. Cell. Neurosci.* **39**, 40-49.
- Yamamori, T. (2011). Selective gene expression in regions of primate neocortex: implications for cortical specialization. *Prog. Neurobiol.* **94**, 201-222.
- Zecevic, N., Chen, Y. and Filipovic, R. (2005). Contributions of cortical subventricular zone to the development of the human cerebral cortex. *J. Comp. Neurol.* **491**, 109-122.